

## APS President Urges Members To Take Action on Federal Science Funding

By Calla Cofield

In response to the FY08 budget passed by Congress, which fell nearly a billion dollars short in science funding compared to the levels authorized last summer, APS president Arthur Bienenstock sent two email messages to all APS members, urging them to write to Congress. The Federal Funding Alert email that went out on January 10 called the cuts a “devastating blow to basic research.” Bienenstock is asking APS members to tell Congress to pass emergency supplemental appropriations to replace some of the cut funding.

The email provided a link to an on-line form for members to write to their representatives and to President Bush, including suggested templates for the letters. At press time, APS’s Washington DC office reported that more than 3300 members had used the online form to write to Congress and the Administration.

The email, titled “Please help rectify science damage in FY08 budget,” emphasized the impact that the budget cuts will have on science programs nationally and internationally:

“The [FY08] budget, which wipes out \$1 billion in increases approved last summer for the National Science Foundation (NSF), the Department of Energy’s Office of Science (DOE Science) and the NIST laboratories, does irreparable damage to science and abandons the Innovation/Competitiveness initiatives of Congress and the Administration.

...The request in the attached let-



Paul Dlugokencky (aDailyCartoon.com) for APS NEWS

ters is to restore that funding in an FY08 supplemental appropriations bill, and to support the FY09 budget at the levels authorized in the COMPETES act, efforts that the APS Washington Office are pursuing with both Congress and the Administration.”

Bienenstock’s email also pointed out that the budget drastically cuts R&D for the International Linear Collider, and zeroes out the US contribution to the ITER project. “These actions are severely damaging to the U.S. standing in the international scientific community,” the message says.

In a second email, sent January 22, Bienenstock added “The Department of Energy Office of Science re-

leased a document last week listing the impacts to all of its programs. In addition to the damage to the Fusion and High Energy Physics programs that I emphasized last week, there are major impacts in Basic Energy Sciences (BES) and Nuclear Physics programs. The Intense Pulsed Neutron Source at Argonne National Laboratory is being closed permanently and various construction projects will be delayed...nearly 700 proposals responding to a BES solicitation for energy research have been declined.”

The online form to write to Congress can be found on the APS website. Go to the Policy and Advocacy page, click on “Advocacy Tools” and then on “Write Congress.”

## POPA’s Short Reports Give Congress Timely Scientific Expertise

The APS Panel on Public Affairs (POPA) has produced a series of “short reports” on topics ranging from energy and the environment to national security issues since 2004. The aim: to provide critical technical expertise to Congress in a timely fashion on policy issues with a strong science or technological component.

The APS membership includes eminent physicists with expertise that is highly relevant to several issues being debated in Congress, and part of POPA’s mission is to provide the Society’s input on those issues to legislators who are responsible for making policy decisions. One way of doing this is through in-depth technical studies, known as full-or short-length reports.

Full-length reports—such as the landmark 1989 Directed Energy Weapons study—are costly and can take as long as three years to complete. By the time a full-length study is completed, Congress may

have acted on some of the pending questions—without input from key scientific experts. Short reports are designed to fill that gap. They can run about 20 pages, include a summary of the main findings and recommendations, and can be completed in eight months or less.

POPA members propose topics for short reports to the panel, which discusses the merits, time scale, logistical feasibility and “whether or not the physics community has something intelligent to say about the issue,” says Robert A. Eisenstein, who chaired POPA in 2007.

“This is very critical,” he says. “We don’t get involved with issues where we don’t have expertise. Our focus is, what can science tell you? We stay out of the political dimension.” If the proposed topic passes muster, a formal charge is prepared.

“The model has worked well because the issues are still fresh in Congress’s mind when we an-

nounce our results,” says Francis Slakey, associate director of public affairs in the APS Washington Office.

The office now receives direct queries from congressional and federal offices on specific issues because staffers are aware that APS has expertise and can respond in a shorter time frame. For example, the U.S. Department of Homeland Security recently asked POPA to convene a panel of experts to evaluate the capability of devices to detect nuclear materials and/or radiation shielding.

Most importantly, Congress seems to be open to the physics community’s recommendations. The first short report—on the planned \$1.2 billion Hydrogen Initiative—appeared in 2004, calling for a focus on basic research and away from demonstration projects. It concluded that major scientific breakthroughs are needed to make

POPA continued on page 11

## 期刊将用中文，日文和朝鲜文印刷作者名字

The APS journals now offer authors the option to include their names in either Chinese, Japanese or Korean characters following the name as it appears in Latin characters, for example: Tadanori Minamisono (南園 忠則) or Chang Kee Jung (정창기). The program, announced in December, is offered for author by-lines throughout the *Physical Review* journals, including *Physical Review Letters*.

The option offers advantages to these authors and to readers of the journal. Many names that are different when expressed in characters become the same when transliterated into English. Showing the characters after the transliterated name removes the ambiguity, and enables readers to know definitively whose work is whose.

The program is the brainchild of Gene Sprouse, APS Editor-in-Chief.

“A person’s name is important. It is the first word that a child learns to write, and it stays with him or her throughout life,” he observed. “Authors who choose to have their names printed this way can show their name on their paper to a friend or family member who may not read English! Our international submissions are growing and we occasionally have trouble ourselves distinguishing one Asian author from another. We value these authors and we want to be welcoming to them in our journals,” he said.

The pilot program for Chinese, Japanese and Korean characters is now available, and with time and experience additional languages may be offered. Instructions for authors on how to supply the proper Unicode characters at the time of submission are at <http://authors.aps.org/names.html>.

## High-Energy Labs Reel Under Budget Cuts

By Ernie Tretkoff

Last year both the Administration and Congress had shown support for increasing spending on physical science; the bipartisan America COMPETES bill authorized significant increases for basic science. But in December, Congress, scrambling to pass an omnibus appropriations bill for fiscal year 2008 that would meet the President’s spending target, cut billions of dollars, including substantial cuts for science.

The impact on science was very broad. At the DOE Office of Science, fusion energy sciences was 33% below the President’s request (including the cancellation of the promised \$160 million US contribution to ITER, an international fusion program); basic energy sciences was 15.3% below; and nuclear physics 8.2% below.

The budget for high energy physics was slashed 12%, from the \$782 million requested to \$688 million.

The cuts will result in layoffs of hundreds of workers at both SLAC and Fermilab. At SLAC, the B-factory experiment will end in March, seven months before its planned shutdown.

In the final appropriations, Fermilab’s budget for FY08 was cut from the \$372 million requested to \$320 million. This is less than the FY07 budget of \$342 million.

The cuts will result in layoffs of about 200 of the lab’s approximately 2000 staff members, and remaining staff will subject to a “rolling furlough,” requiring them to take two to three days of unpaid leave per month. Work on development for future projects, such as the ILC, has been stopped at Fermilab.

Fermilab Director Pier Oddone called the budget tremendously disruptive. He said it pained him to have implement layoffs and furloughs, but says they are necessary to keep the

LABS continued on page 4

## Abstract Reasoning



Photo by Ken Cole

Marco Fornari of Central Michigan University and Noam Bernstein of the Naval Research Laboratory were among the 132 APS members who came to College Park, MD in December to sort the almost 7000 abstracts that had been submitted to the March Meeting. The meeting takes place in New Orleans, March 10-14.

## Members in the Media



“We would probably support any competent scientist that wants to run for Congress.”

**Leon Lederman**, on getting scientists to run for office, *US News and World Report*, December 6, 2007

“This represents an extraordinary waste of the investment and leadership established by the UK in this truly international project.”

**Albrecht Wagner**, *DESY*, on the UK pulling out of the ILC, *The Telegraph* (UK), December 13, 2007

“It reminds me a little bit of NASA’s decision to launch the space shuttle with O ring problems.”

**Nigel Lockyer**, *TRIUMF*, on Canadian Prime Minister Harper’s decision to overrule Canada’s nuclear safety regulators and fire up the Chalk River reactor, *Vancouver Sun*, December 13, 2007

“Can the models accurately explain the climate from the recent past? It seems that the answer is no.”

**David H. Douglass**, *University of Rochester*, on climate models, *Fox News.com*, December 13, 2007

“If there’s a math of knots there should also be a science of knots,

something that explains why they form.”

**Douglas Smith**, *UC San Diego*, *San Diego Union Tribune*, December 19, 2007

“He was a real nuisance when I was taking high school physics. It was the classic thing. . . you ask for help and all you really want to know is the answer to problem 5B—and he wants to explain it to you.”

**Persis Drell**, *SLAC*, on being the daughter of a physicist, *San Jose Mercury News*, December 18, 2007

“If you tell 100 million people to go east, 25 million will go west because they don’t trust the government.”

**Jay C. Davis**, on communication after a nuclear attack, *Los Angeles Times*, January 6, 2008

“Ours is a small detector. It’s roughly the size of this building.”

**Brad Cox**, *University of Virginia*, on the CMS detector being built for the LHC, *Richmond Times-Dispatch*, January 9, 2008

“When you break an egg and scramble it you are doing cosmology.”

**Sean Carroll**, *Catech*, *The New York Times*, January 15, 2008

## Learning Assistants Impact Undergraduate Teaching

At a recitation section for an introductory physics course at the University of Colorado at Boulder, groups of students are typically seated around tables discussing the nuances of Newtonian mechanics or the intricacies of electromagnetic induction—but it’s not easy to spot the TA. In fact, there are probably several assistants in the room, but they don’t lecture or work problems on the board. Instead, they move from table to table asking students questions to elicit their misconceptions about physics, and guiding them toward a more sophisticated understanding of the concepts being taught. These undergraduate Learning Assistants not only help their peers learn physics, but often discover a passion for teaching in the process.

Now four years old, the Colorado Learning Assistant program is starting to attract attention from science faculty around the country who want to recruit strong students into teaching careers. To capitalize on this momentum, the Physics Teacher Education Coalition (PTEC)—a project led by APS in collaboration with AAPT

and AIP—recently organized and sponsored a two-day workshop that brought 22 faculty members from a diverse group of 14 colleges and universities to Boulder to learn how to replicate Colorado’s successes on their own campuses.

The workshop, which was led by a team of Colorado science and education professors, caught the Learning Assistants in action both in the recitation sessions described above, and in the weekly science pedagogy course that all Learning Assistants take during their first year in the program. This course provides the crucial opportunity for the program leaders to impress upon the young teachers the importance, and difficulty, of truly engaging students. As Valerie Otero, an education professor who co-teaches the class, put it, “the Learning Assistant experience helps students realize that teaching is a real intellectual challenge, and for many of them, this is exactly what they’re looking for.” To demonstrate this, she had workshop participants and Learning Assistants

**ASSISTANTS continued on page 3**

## This Month in Physics History

### Heisenberg and the Uncertainty Principle

In February 1927, the young Werner Heisenberg developed a key piece of quantum theory, the uncertainty principle, with profound implications.

Werner Heisenberg was born in December 1901 in Germany, into an upper-middle-class academic family. He liked mathematics and technical gadgets as a boy, and his teachers considered him gifted. In 1920 he began studies at the University of Munich, and published four physics papers within two years under the guidance of mentor Arnold Sommerfeld. Heisenberg became professional friends with Wolfgang Pauli, who was just one year older than Heisenberg and also a student at Munich.

He earned his doctorate in 1923, with a thesis on a problem in hydrodynamics, though he nearly failed due to his poor performance on the required experimental questions on the oral examination. After receiving his doctorate, he worked as an assistant to Max Born at Göttingen, then spent a year working with Niels Bohr at his institute in Copenhagen.

The prevailing quantum theory in the early 1920s modeled the atom as having electrons in fixed quantized orbits around a nucleus. Electrons could move to higher or lower energy by absorbing or emitting a photon of the right wavelength. The model worked well for hydrogen, but ran into problems with larger atoms and with molecules. Physicists realized a new theory was necessary.

Heisenberg objected to the current model because he claimed that since one couldn’t actually observe the orbit of electrons around a nucleus, such orbits couldn’t really be said to exist. One could only observe the spectrum of light emitted or absorbed by atoms. Starting in 1925, Heisenberg set to work trying to come up with a quantum mechanics that relied only on properties that could, at least in theory, be observed.

With help and inspiration from several colleagues, Heisenberg developed a new approach to quantum mechanics. Basically, he took quantities such as position and velocity, and found a new way to represent and manipulate them. Max Born identified the strange math in Heisenberg’s method as matrices. The new formulation accounted for many observed properties of atoms.

Shortly after Heisenberg came up with his matrix-based quantum mechanics, Erwin Schrödinger developed his wave formulation. The absolute square of Schrödinger’s wave function was soon interpreted as the probability of finding a particle in a certain state. Schrödinger’s wave formulation, which he soon proved was mathematically equivalent to Heisenberg’s matrix methods, became the more popular approach, partly because physicists were more comfortable with it than with the unfamiliar matrix mathematics. The unpopularity of his own method annoyed Heisenberg, especially because a lot was at stake at the time as he and other young scientists were beginning to look for their first jobs as professors as an older generation of sci-

entists was retiring.

Though others may have found the wave approach easier to use, Heisenberg’s matrix mechanics led him naturally to the uncertainty principle for which he is well known. In matrix mathematics, it is not always the case that  $axb = bxa$ , and for pairs of variables that don’t commute, such as position and momentum, or energy and time, an uncertainty relation arises.

Heisenberg conducted a thought experiment as well. He considered trying to measure the position of an electron with a gamma ray microscope. The high-energy photon used to illuminate the electron would give it a kick, changing its momentum in an uncertain way. A higher resolution microscope would require higher energy light, giving an even bigger kick to the electron. The more precisely one tried to measure the position, the more uncertain the momentum would become, and vice versa, Heisenberg reasoned. This uncertainty is a fundamental feature of quantum mechanics, not a limitation of any particular experimental apparatus.

Heisenberg outlined his new principle in 14-page a letter to Wolfgang Pauli, sent February 23, 1927. In March he submitted his paper on the uncertainty principle for publication.

Niels Bohr pointed out some errors in Heisenberg’s thought experiment, but agreed the uncertainty principle itself was correct, and the paper was published.

The new principle had deep implications. Before, it had been thought that if you knew the exact position and momentum of a particle at any given time, and all the forces acting on it, you could, at least in theory, predict its position and momentum at any time in the future. Heisenberg had found that not to be true, because you could never actually know a particle’s exact position and momentum at the same time.

The uncertainty principle soon became part of the basis for the widely accepted Copenhagen interpretation of quantum mechanics, and at the Solvay conference in Brussels that fall, Heisenberg and Max Born declared the quantum revolution complete.

In the fall of 1927, Heisenberg took a position as a professor at the University of Leipzig, making him the youngest full professor in Germany. In 1932 he won the Nobel Prize for his work on quantum mechanics. He continued his scientific research in Germany. During World War II, though he was not a member of the Nazi party, he was a patriotic German citizen, and he became a leader in the German fission program, which failed in its effort to build an atomic bomb. Heisenberg’s actions and motivations have been the subject of controversy ever since. He died in 1976.

Reference/further reading: David Cassidy, *Uncertainty: the Life and Science of Werner Heisenberg* (New York: W.H. Freeman, 1992).



## APS NEWS

Series II, Vol. 17, No. 02  
February 2008

© 2008 The American Physical Society

Coden: ANWSEN ISSN: 1058-8132

Editor . . . . . Alan Chodos  
Staff Writer . . . . . Ernie Tretkoff  
Contributing Editor . . . . . Jennifer Ouellette  
Art Director and Special Publications Manager . . . . . Kerry G. Johnson  
Design and Production . . . . . Nancy Bennett-Karasik  
Proofreader . . . . . Edward Lee

APS News (ISSN: 1058-8132) is published 11X yearly, monthly, except the August/September issue, by the American Physical Society, One Physics Ellipse, College Park, MD 20740-3844, (301) 209-3200. It contains news of the Society and of its Divisions, Topical Groups, Sections and Forums; advance information on meetings of the Society; and reports to the Society by its committees and task forces, as well as opinions.

Letters to the editor are welcomed from the membership. Letters must be signed and should include an address and daytime telephone number. The APS reserves the right to select and to edit for length or clarity. All correspondence regarding APS News should be direct-

ed to: Editor, APS News, One Physics Ellipse, College Park, MD 20740-3844, E-mail: letters@aps.org.

**Subscriptions:** APS News is an on-membership publication delivered by Periodical Mail. Members residing abroad may receive airfreight delivery for a fee of \$15. Nonmembers: Subscription rates are available at <http://librarians.aps.org/institutional.html>.

**Subscription orders, renewals and address changes** should be addressed as follows: For APS Members—Membership Department, American Physical Society, One Physics Ellipse, College Park, MD 20740-3844, membership@aps.org.

For Nonmembers—Circulation and Fulfillment Division, American Institute of Physics, Suite 1N01, 2 Huntington Quadrangle, Melville, NY 11747-4502. Allow at least 6 weeks advance notice. For address changes, please send both the old and new addresses, and, if possible, include a mailing label from a recent issue. Requests from subscribers for missing issues will be honored without charge only if received within 6 months of the issue’s actual date of publication. Periodical Postage Paid at College Park, MD and at additional mailing offices. Postmaster: Send address changes to APS News, Membership Department, American Physical Society, One Physics Ellipse, College Park, MD 20740-3844.

#### APS COUNCIL 2008

**President**  
Arthur Bienenstock\*, *Stanford University*  
**President-Elect**  
Cherry Murray\*, *Lawrence Livermore National Laboratory*  
**Vice-President**  
Curtis G. Callan, Jr.\*, *Princeton University*  
**Executive Officer**  
Judy R. Franz\*, *University of Alabama, Huntsville (on leave)*  
**Treasurer**  
Joseph W. Serene\*, *Georgetown University (emeritus)*

#### Editor-in-Chief

Gene Sprouse\*, *Stony Brook University (on leave)*

#### Past-President

Leo P. Kadanoff\*, *University of Chicago*

#### General Councillors

Robert Austin, Christina Back\*, Marcela Carena, Elizabeth Beise, Katherine Freese, Wendell Hill\*, Ann Orel\*, Richard Slusher\*

#### International Councillor

Sabayasachi Bhattacharya

#### Chair, Nominating Committee

Philip Phillips

#### Chair, Panel on Public Affairs

Miles Klein

#### Division, Forum and Section Councillors

Charles Dermer (*Astrophysics*), P. Julienne (*Atomic, Molecular & Optical Physics*) Robert Eisenberg (*Biological*), Charles S. Parmenter (*Chemical*), Arthur Epstein (*Condensed Matter Physics*), (Computational-TBA), James Brasseur (*Fluid Dynamics*), Peter Zimmerman\* (*Forum on Education*), Roger Stuewer (*Forum on History of Physics*), Stefan Zollner (*Forum on Industrial and Applied Physics*), David Ernst\* (*Forum on International Physics*), (*Forum on Physics and Society-TBA*), Steven Rolston (*Laser Science*), Leonard Feldman\* (*Materials*), Akif Balantekin\* (*Nuclear*), Janet Conrad (*Particles & Fields*), Ronald Ruth (*Physics of Beams*), David Hammer (*Plasma*), Scott Milner (*Polymer Physics*), Paul

Wolf (*Ohio Section*), Heather Galloway (*Texas Section*)

#### ADVISORS

**Representatives from Other Societies**

Fred Dylla, AIP; Lila Adair, AAPT

#### International Advisors

Francisco Ramos Gómez, *Mexican Physical Society*  
Louis Marchildon, *Canadian Association of Physicists*

#### Staff Representatives

Alan Chodos, *Associate Executive Officer*; Amy Flatten *Director of International Affairs*; Ted Hodapp, *Director of Education and Diversity*; Michael Lubell, *Director, Public Affairs*; Dan Kulp, *Editorial Director*; Christine Giaccone, *Director, Journal Operations*; Michael Stephens, *Controller and Assistant Treasurer*

#### Administrator for Governing Committees

Ken Cole

#### \* Members of the APS Executive Board

# Profiles in Versatility

## Physics Major Facilitates Success in Speechwriting (and the Funny Business)

By Alaina G. Levine

There is professional lore that claims a person changes careers on average seven times in their lives. Michael Long may demonstrate an element of truth to this. By age 31, he had already held positions in physics, comedy, mathematical modeling, and freelance writing, and by no means had he reached his peak yet. Only months after his 32<sup>nd</sup> birthday, the physics aficionado entered the ring of political speechwriting after being hired by then Tennessee Senator Fred Thompson.

"Senator Thompson was a rising star in politics and I was thrilled to start in politics with such an exciting opportunity," Long recalls.

Today, Long is a well-known writer of speeches (and other pieces), a line of work he equates to being "a professional explainer".

"I look at a complicated subject and make it understandable and interesting to a lay audience," he states. Long especially thrives with matters of technology, healthcare, and domestic policy.

Nowadays, he does mostly freelance speeches, although he also serves as a director of the White House Writers Group, a communications firm specializing in business and public policy, through which he has authored and ghost-authored numerous speeches, op-eds and white papers for some of the country's most well-known politicians, including President George W. Bush.

To the amateur, the speechwriting process seems similar to building an early-stage technology enterprise. It is characterized by extensive research and development (in this case of words and ideas) that must eventually lead to a product (the speech) a customer will buy. This is where physics expertise comes in handy.

Long uses his scientific background as a springboard to organize his thoughts and analyze what words to choose.

However, ultimately his success at both writing and attracting and keeping clients is dependent on a "dirty secret" of the field: "most clients want to sound like themselves, only better," Long reveals. So when a client requests that Long writes a speech in the client's "own voice", "if it is articulate and well-written, whatever you give someone they think it's in their voice," he says.

Long received his BS in physics from Murray State University, and remembers that he was initially drawn to the subject because "I wasn't concerned about impressing anybody," he says. Physics "seemed new to me. I didn't know any physicists; it was like someone from the moon. But it's a wonderful way to think."

He enrolled in the graduate program at Vanderbilt University because "the idea of finding order in disorder appealed to me," he says. "Physics teaches you to approach things looking for patterns and processes." However, like many physics students, Long came to the realization that "I was not going to be the king of the mountain," he admits. "In my professional life, whatever I did I wanted to be the very best at it, and when I got into grad school, physics did not come as naturally to me as other things. I saw so many people that were hard-wired to explore new knowledge... I didn't think I had their skills."

Long recognized "If I want to make a difference, this isn't going to be where it's going to happen."

To make such a difference, Long took what many would consider a long-shot: he left graduate school to

pursue a full-time career as a comedian. And for a while he did pretty well, including an invitation to be the house emcee at a prestigious comedy club in Nashville.

Unfortunately, jobs in comedy, much like physics, are difficult to come by and often do not pay well (although on the bright side, comedians don't have to deal with the promotion and tenure process). So he tapped into his technical background to pay the bills and got a job



at a software company where he did mathematical and systems modeling and software design for the telecommunications industry. He stuck with comedy, however, regarding it at the time as an avocation.

But the life of a comedian involves constant writing, and Long developed a fondness and talent for putting his words to paper. He embarked on a career writing articles and op-eds. He quickly found success and was published in local and national publications.

Undeniably a devotee to career diversity, Long leveraged his writing projects into yet another line of busi-

ness in political speechwriting. He had always had a penchant for politics, and in 1994, began researching the subject. He read political articles and noticed that many of the authors were speechwriters. "They had backgrounds like mine," he says. "None of these writers were journalists."

Long realized that even though he had not taken an English class since high school, his varied, scientifically-based skills would help him to be not only a great writer, but a great speechwriter as well. After all, unlike other majors, "I had studied physics," something that is "freaking useful," he laughs. He now advises students that "if you study physics you will learn something not many people know: you will learn how to think, and you will be the most thoughtful, critical writer to ever pick up a pen."

He commenced on his new career by cold-writing a letter to Senator Thompson. Long explained to the politician that "I am not a speechwriter but I want to be one and I want to be yours." He sent him clips of his work and soon was hired as the sole speechwriter for Thompson.

In addition to providing skills such as general problem-solving, physics has done something especially valuable for Long. "It's an impressive credential and it acts as a talisman: when you are going to talk about something technical, people tend to defer to you," he says. "There's a mystique and myth around it, and people are impressed by it, so they give you a little more latitude to offer an opinion on pretty much anything."

Although speechwriting is his current passion and main source of income, Long has amassed a very long list of other, very unusual accomplishments. He has written a syndicated newspaper column, and

the liner notes for the DVD of Jerry Seinfeld's movie "Comedian". He is a guest host on the nationwide Radio America network, has taught classes in stand-up comedy, and was a Comic Relief competition winner. He is also a consultant for the American Film Renaissance, a non-profit film organization, and has served as a consultant for material for Saturday Night Live and in *The Onion*, an internationally-known satirical newspaper and website.

Long has an impressive list of contacts in the comedy field. He counts comedian (and former Nixon speechwriter) Ben Stein as a friend, who observes that Long is "a poet... I don't see anyone out there under the age of 50 writing better short pieces than he is right now."

Now in his early 40s, Long is looking to add yet more diversity to his career trajectories. In November 2007, he took a mere 30 days to write his second novel. And last month he began teaching public relations writing at the graduate school at Georgetown University. He sees physics as an asset and in harmony with his professional goals and strategies.

"What I do is what a physicist does, only carried to another realm: stripping the superfluous material to its core and describing it as what it is," Long says. "[Physics] is what makes me an unusual writer and a writer who can make a living at writing. I look for the elements of the problem, throw away the other stuff and pass the piece along all shiny and polished up so people can understand it."

Copyright, 2007, Alaina G. Levine.

Alaina G. Levine can be contacted through her website [www.alaina-levine.com](http://www.alaina-levine.com).

### ASSISTANTS continued from page 2

work together to figure out how to guide hypothetical non-science majors from a naïve conceptual model of a phenomenon—in this case magnetization—toward a more sophisticated model accepted by scientists.

Other workshop sessions focused on the importance of collecting data to measure the impact of the program. Physics professors Steve Pollock and Noah Finkelstein have amassed years' worth of assessment results that demonstrate that the program has improved the conceptual understanding of students in introductory physics and of the Learning Assistants themselves. They also pointed out that at just \$1,500 a semester, a department can hire numerous Learning Assistants for the price of one graduate TA.

But perhaps the program's greatest success has been as a teacher recruitment tool. Nine Learning Assistants were enrolled in physics and astrophysics teacher certification programs at Boulder in 2005-2006, as compared to one student

(who was not a Learning Assistant) in 2004-2005, when only five students in the entire state of Colorado were enrolled in physics and astrophysics teacher certification programs. Ted Hodapp, Director of Education and Diversity at APS, explained the program's appeal. "Learning Assistant programs are a kind of surprise attack for students who haven't previously thought of themselves as potential teachers. By giving students a low-stress, low-commitment early teaching experience, physics departments can get them excited about teaching and recruit them into teacher preparation programs."

The Colorado program turns this excitement into a career track by requiring Learning Assistants who want to remain in the program after two semesters to enter a teacher certification program. This makes them eligible for NSF-funded Noyce Teaching Fellowships of up to \$10,000 a year, in exchange for a commitment to teach in a high-needs school after graduation.

Noyce Fellows also get invited to serve as mentors for novice Learning Assistants, and collaborate with faculty on education research projects. Otero strongly encouraged workshop participants to provide this money for their future teachers.

At a question-and-answer session during the workshop, several experienced Learning Assistants shared their perspectives on the program. They spoke about the culture change that the program has catalyzed within science departments, as students and some professors have begun to view teaching as a respectable and worthwhile career for a science major to pursue. They also noted the tight learning community that develops among Learning Assistants, many of whom remain in the program for multiple semesters. Most workshop participants indicated that they appreciated the opportunities to interact with real, live Learning Assistants.

Most faculty attendees described specific plans they had generated

during the workshop to develop elements of the program on their campuses. To sustain and spread the enthusiasm that the workshop generated, PTEC is developing a web-

based Learning Assistant "home" (<http://www.ptec.org/conferences/CULA/>), and hopes to organize future conference sessions on Learning Assistant programs.



Photo by Monica Plisch/APS Staff

Learning Assistant Allison Lanini and Colorado Education Professor Valerie Otero collaborate on a demonstration of interactive teaching.

# Letters

## Ethicist Gets It Right

I loved your "Ask the Ethicist" column in the December *APS News*. I recall having to write almost all of my own recommend letters for grad school from my profs and later for employment as well. Now that roles are reversed, I always require an electronic copy (for me to edit as

appropriate) of a recommend letter be provided to me from the requestor and assume my peers and colleagues expect the equivalent. Very nice synopsis from Jordan Moiers.

**Robert B. Hayes**  
*Las Vegas, NV*

## Ethicist Gets It Wrong

Unbelievable! A student writes in asking if it's ethical to ghost write his/her own letter of recommendation, and there is no outrage that the person (most likely a professor) asked for this ghost written letter in the first place... THE EMPEROR IS NOT WEARING ANY CLOTHES! How about this: if the professor really wants to get to know the student in order to write a good letter of recommendation, they sit down with the student for an hour or so and talk to them. Then, the professor could take another hour to craft a letter that captures his impression of the student based on interactions in class and the personal discussion. Unfortunately, that would take at a minimum two hours of time away from the professor's very busy

schedule, and in my opinion this is the heart of the professor's request. Let's not delude ourselves; the professor is not going to modify the letter. The professor is going to read the letter over quickly, and so long as it sounds reasonable is going to sign it and get that nuisance-task off his or her plate. Interacting with students is not a top priority for a number of faculty, especially at the large research institutions, no matter what lip service they pay to it. Shame on the professor for asking for this ghost written letter, and shame on "the Ethicist" for not calling the professor to task.

**James Camparo**  
*Redondo Beach, CA*

## Bias? What Bias?

I do not believe that you have fallen into "American Olympic athletes" trap, as suggested in the Editor's Note to the letter from Bob Dewar in the January *APS News*. After all, APS means American Physical Society, so its focus is naturally American Physics. If Bob Dewar needs information about Australian Olympians I am sure that there exists a relevant Australian periodical. I do not advocate limiting news to exclusively American, but I do advocate a reasonable balance, and as someone who is not an Ameri-

can (I am Polish), I have no bias on the "America versus the Rest of the World" issue.

**Piotr Zolnierczuk**  
*Bloomington, IN*

**Ed. Note:** We thank Piotr Zolnierczuk for his support. We are proud of the roughly 20% of APS members who are not in the US, and we attempt, in part, to reflect that constituency with our "International News" column that appears every other month.

## LABS continued from page 1

lab running. "I get paid to optimize the program, not to protect every last job," he said. He said the most serious impact of the cuts was the effect on the lab's future, as projects that would be central to Fermilab's future program have had funding cut off. "We're not getting ready for what happens 2-3 years down the line," said Oddone. "That's the biggest impact."

Fermilab, the nation's only laboratory devoted solely to high energy physics, is especially vulnerable to budget cuts, Oddone pointed out, but he does not plan to change the lab's mission. Oddone said he is waiting for the release of the President's FY09 budget in early February before making further decisions.

Several Fermilab scientists said they felt that Congress had overlooked the serious consequences for physics in their haste to pass the enormous omnibus appropriations bill. They said that this budget sends a bad message to young people, who might be deterred from going into physics, and suggests that the US is not a reliable partner for large international endeavors.

The \$60 million expected for FY08 for ILC development was reduced to \$15 million, but since the cuts were made in December, three months into the 2008 fiscal year, that money has already been spent. "It's effectively stopped all work," said Fermilab's ILC program director Robert Kephart.

The budget situation puts Fermilab

in jeopardy as the site for the ILC. "If we don't reform the budget process, the world will want to build the ILC elsewhere," said Oddone.

"It makes the US look like a poor partner," for international collaboration, said Kephart.

The United Kingdom has also withdrawn its promised contribution to the ILC.

Large particle physics experiments such as the ILC need to be planned years in advance. "The damage from this kind of budget process is really severe," he said.

Funding has also been eliminated for an experiment called NOvA, which would have become the main neutrino program at Fermilab and was about to begin construction. NOvA was designed to search for muon neutrino to electron neutrino oscillations, a key to unanswered questions in neutrino physics. About \$36 million had been expected for the project. "It's a difficult situation. You can't just take a project like this and stop now and start later," said Gary Feldman, a spokesman for NOvA. It is unclear how long NOvA will be delayed. "Right now there's a lot of uncertainty," said Mark Messier, another physicist working on NOvA. "The message it sends to graduate students is terrible," he said.

While work on future projects has stopped, Fermilab's Tevatron will continue to run until 2009 as planned. Spokesmen for the CDF and DZero, Fermilab's two main detector collaborations, said the Tevatron could

## Oil Addiction Distorts Senator's Thinking

Senator Byron Dorgan's Back Page [*APS News*, December 2007] fell short of the frankness and completeness I was hoping for in dealing with America's oil addiction.

Most glaring was Senator Dorgan's statement of "...so they don't disrupt our energy supplies." It is not our energy. It is theirs! The Senator states unequivocally that America needs to "improve management of alliances to better secure global oil supplies." This thinking is equivalent to China arguing for international pressure to ensure "their" corn, which happens to be growing in Kansas, is kept inexpensive so it won't disrupt China's economy. Our addiction to oil is distorting our philosophy for the worse.

Dorgan is arguing two main points: (1) increased energy conservation/alternative energy development, and (2) greater production of oil at home and strengthened alliances so import prices remain low. These two points are like arguing for both n and 1/n

to get larger as n goes to infinity. To be frank, the American energy consumer will not conserve or arduously look for alternative sources so long as oil supply is plentiful which, in turn, keeps prices low.

By plotting the percent of energy consumption in the US that comes from renewable resources as a function of time, one finds the graph is flat from 2001 to 2005. It has stayed at about 6% (or 3% if one removes ethanol from a renewable classification). A sustainable, responsible, and secure future cannot be had until this graph shows a significant, positive trend.

Senator Dorgan did little to belie the false impression that corn-based ethanol is a panacea for energy independence and good for the environment. The explicit support for increasing "renewable fuels like ethanol" is disappointing.

Making progress with CAFÉ standards is good but will soon be negated by increased number of vehicles on the road. Why is the fundamental

problem of population pressure on natural resources so absent from our political discourse? Senator Dorgan implicitly believes—along with many politicians from the President on down—that money generated by economic growth can be applied to fix any problem growth creates. This is just plain pollyannaish. One can argue economic growth does not necessarily imply population growth but disconnecting the two takes immense strain.

**Alan J. Scott**  
*Menomonie, WI*

## Nuclear Energy Too Controversial?

An entire page on energy security and methods to achieve it, without a single mention of nuclear energy. That must exceed Senator Dorgan's tolerance for controversial issues.

**John Tanner**  
*Idaho Falls, ID*

## US Needs a Tax on Oil

The Back Page article by Senator Dorgan in the December *APS News* is absolutely correct about the urgent need to find new, renewable, and environmentally friendly energy sources, and to conserve our energy so we need less of it. But I am disappointed by the lack of political will to do the most effective thing. A tax on

oil provides a meaningful incentive to conserve, and makes alternative energy a better investment. An oil tax can also raise revenue to fund research in renewable energy. I favor a floor price for oil, so investors could be assured their investment in alternate energy will not go bust. Almost all the industrialized nations in the world except

the United States tax oil to encourage conservation and investment in oil alternatives, and it is time that the United States developed the political will to do the same thing.

**Paul McManamon**  
*Dayton, OH*

## Hubble's Thoughts on Hubble's Law

I always enjoy reading "This Month in Physics History" and the January installment on Hubble's discoveries was no exception. However, I would like to point out a few minor errors in that piece. Most astronomers in the early 20s favored the theory that spiral nebulae were "island universes" and in fact believed the Milky Way to be much smaller than we now know it to be. Shapley and a few others favored the idea of a much larger Milky Way which contained the spiral nebulae, but Shapley's letters indicate that he knew he was in the minority on this issue. Also, it was Henry Norris

Russell who presented (on behalf of Hubble) the data on Cepheids in Andromeda at the AAS meeting in January 1925. Most importantly, it is untrue that "Hubble didn't discuss the implications of what he had found" in his 1929 PNAS paper. In the final paragraph of that paper he writes "the velocity-distance relation may represent the de Sitter effect," referring to the model of the Universe presented by Willem de Sitter in 1917. This model was originally interpreted as a static model, but did predict a redshift that increased with distance because of scattering and an apparent slowing down of distant atomic vibrations.

So in 1929 Hubble did not interpret his data as indicating an expanding Universe, but rather as supporting de Sitter's static model. It was only later realized that de Sitter's model was equivalent via a coordinate transformation to expanding models such as that proposed by Georges Lemaître in 1927 (Lemaître's model was unknown to Hubble and most astronomers until 1930). A detailed account of this history is given in Robert W. Smith's *The Expanding Universe* (Cambridge U Press, 1982).

**Todd Timberlake**  
*Mount Berry, GA*

still do exciting science. They pointed to many recent accomplishments of the Tevatron, including precision measurements of the mass of the top quark and W and Z bosons, discovery of some new particles and detection of some rare processes such as the production of single top quarks. CDF spokesman Jacobo Konigsberg of the University of Florida said that the Higgs particle mass is probably within the energy range accessible to the Tevatron, and finding the Higgs is a matter of collecting enough data before the machine shuts down. There is also the possibility of finding new physics. "Unless you search, you cannot find," said Fermilab scientist and DZero spokesman Dmitri Denisov.

Although they are excited about the prospect of new discoveries from the Tevatron, Fermilab scientists worried about the future of US particle physics. Young people will see the uncertainty in funding and will not be attracted to the field, and people affected by layoffs generally do not return to the field. "If people see no future, they will try to look for something else" said Denisov.

These budget cuts could be detrimental not only to high energy physics, but to the broader economy,

because these accelerators advance technology, and students who work on these projects gain a wide range of valuable skills, including communications skills, computing skills and experience with electronics and cutting edge technology. "There are very few endeavors as varied and rich as high-energy physics," said Robert Roser, a Fermilab scientist and CDF spokesman.

Fermilab is also continuing its contributions to the Large Hadron Collider at CERN. Dan Green, Fermilab's spokesman for CMS, one of the LHC detectors, said he was excited about work on CMS, much of which can be done in the United States. The CMS detectors are being tested, and are on schedule to be ready before the LHC begins operation. "LHC is our only frontier device now," said Green. Many US scientists are participating in that collaboration, and the US remains committed to the project. "We have been good international partners," he said, but the recent budget cuts are worrisome.

While the future is uncertain, Fermilab officials and scientists say they are hopeful that this year's budget is a one-time setback from which the lab could recover. But some are less opti-

mistic about future funding for physics. SLAC Director Persis Drell told an all hands meeting at SLAC in January, "While the bipartisan enthusiasm for the physical sciences appears to be strong and will likely continue, we cannot view the current budget challenge as a temporary setback. I do not believe that the physical science budgets will grow as quickly as has been hoped for...I believe that nationally we will need to adjust to a smaller base program going forward."

Officials in towns near Fermilab have passed resolutions urging Congress to restore funding to the lab, and Illinois Senators Dick Durbin and Barack Obama and Congresswoman Judy Biggert have sent letters to Jim Nussle, Director of the Office of Management and Budget, asking for increased funding for high energy physics in FY09.

"I really feel grateful for all the support we're getting," said Fermilab Deputy Director Young-Kee Kim. She stressed the need for scientists to communicate with the public and with lawmakers so they understand the value of basic science.

# Physics News in 2007

A Supplement to APS News

Edited by Phil Schewe, Ben Stein and Jason Bardi

## Introduction

*Physics News in 2007*, a summary of physics highlights for the past year, was compiled from items appearing in AIP's weekly newsletter *Physics News Update*, written by Phil Schewe, Ben Stein and Jason Bardi. (Ben Stein has since left AIP and is now at NIST. Jason Bardi has replaced Ben Stein at AIP)

The items below are in no particular order. Because of limited space in this supplement, some physics fields and certain contributions to particular research areas might be under-represented in this compendium. These items mostly appear as they did during the year, and the events reported therein may in some cases have been overtaken by newer results and newer publications which might not be reflected in the reporting. Readers can get a fuller account of the year's achievements by going to the *Physics News Update* website at <http://www.aip.org/pnu> and APS's *Physical Review Focus* website at <http://focus.aps.org/>.

## Gravitational Wave Background

In the standard model of cosmology, the early universe underwent a period of fantastic growth. This inflationary phase, after only a trillionth of a second, concluded with a violent conversion of energy into hot matter and radiation. This "reheating" process also resulted in a flood of gravitational waves.

The gravitational wave background (GWB) dates from the trillionth-of-a-second mark, while the cosmic microwave background (CMB) sets in around 380,000 years later when the first atoms formed. What does the GWB represent? It stems from three different production processes at work in the inflationary era: waves stemming from the inflationary expansion of space itself; waves from the collision of bubble-like clumps of new matter at reheating after inflation; and waves from the turbulent fluid mixing of the early pools of matter and radiation, before equilibrium among them (known as thermalization) had been achieved. The gravity waves would never have been in equilibrium with the matter (since gravity is such a weak force there wouldn't be time to mingle adequately); consequently the GWB will not appear to a viewer now to be at a single overall temperature.

A new paper by Juan Garcia-Bellido and Daniel Figueroa (Universidad Autonoma de Madrid) explains how these separate processes could be detected and differentiated in modern detectors set up to see gravity waves, such as LIGO, LISA, or BBO (Big Bang Observer). First, the GWB would be redshifted, like the CMB. But because of the GWB's earlier provenance, the reshifting would be even more dramatic: the energy of the waves would be downshifted by 24 orders of magnitude. Second, the GWB waves would be distinct from gravity waves from point sources (such as the collision of two black holes) since such an encounter would release waves with a sharper spectral signal. By contrast the GWB from reheating after inflation would have a much broader spectrum, centered around 1 hertz to 1 gigahertz depending on the scale of inflation.

Garcia-Bellido suggests that if a detector like the proposed BBO could disentangle the separate signals of the end-of-inflation GWB, then such a signal could be used as a probe of inflation and could help explore some fundamental issues as matter-antimatter asymmetry, the production of topological defects like cosmic strings, primordial magnetic fields, and possibly superheavy dark matter.

For comparable results see the paper by Easter and Lim in the *Journal of Astroparticle Physics*, JCAP04(2006)010. (Garcia-Bellido and Figueroa, *Phys. Rev. Lett.* **98**, 061302 (2007))

## The Casimir Effect Heats Up

For the first time, a group led by Nobel laureate Eric Cornell at the National Institute of Standards and Technology and the University of Colorado in Boulder has confirmed a 1955 prediction, by physicist Evgeny Lifschitz, that temperature affects the Casimir force, the attraction between two objects when they come to within 5 millionths of a meter of each other or less. These efforts heighten the understanding of the force and enable future experiments to better account for its effects.




Photo by E. Cornell group/JILA

attractive force threefold, confirming theoretical predictions recently made by the group's co-authors in Trento, Italy.

The Casimir force arises from effects of the vacuum. According to quantum mechanics, the vacuum contains fleeting electromagnetic waves, in turn consisting of electric and magnetic fields. The electric fields can slightly rearrange the charge in atoms. Such "polarized" atoms can then feel a force from an electric field. The vacuum's electric fields are altered by the presence of the glass, creating a region of maximum electric field that attracts the atoms. In addition, heat inside the glass also drives the fleeting electromagnetic waves, some of which leak onto the surface as "evanescent waves." These evanescent waves have a maximum electric field on the surface and further attract the atoms.

In their work, the researchers investigated the Casimir-Polder force, the attraction between a neutral atom and a nearby surface. The Colorado group sent ultracold rubidium atoms to within a few microns of a glass surface. Doubling the temperature of the glass to 600 degrees Kelvin while keeping the surroundings near room temperature caused the glass to increase its

Electromagnetic waves from heat in the rest of the environment would usually cancel out the thermal attraction from the glass surface. However, dialing up the temperature on the glass tilts the playing field in favor of glass's thermal force and heightens the attraction between the wall and the atoms. (Obrecht et al., *Phys. Rev. Lett.* **98**, 063201 (2007)) Also see the NIST press release: [http://www.nist.gov/public\\_affairs/newsfromnist\\_casimir-polder.htm](http://www.nist.gov/public_affairs/newsfromnist_casimir-polder.htm))

## Radium Atoms Trapped

Physicists at Argonne National Laboratory, near Chicago, have laser-cooled and trapped radium atoms for the first time.

Surprisingly, room temperature blackbody photons—thermal radiation over a wide spectrum emitted by the apparatus itself—were found to play a critical role in the laser-trapping of this rare and unstable element. This represents the heaviest atom ever trapped by laser light.

Using only 20 nanograms of radium-225 (half-life of 15 days) and one microgram of radium-226 (half-life of 1,600 years), the Argonne scientists held tens of radium-225 and hundreds of radium-226 atoms in the laser trap.

Why go through the trouble of trapping radium atoms? Because it might provide a chance to detect a violation of time-reversal symmetry (abbreviated with the letter T), which would manifest itself as an electric dipole moment in the radium atom.

Electric dipole moment searches have been ongoing for over 50 years and continue to yield smaller and smaller limits on the size of these T-violating interactions. These limits place constraints on theories beyond the Standard Model of particle physics and explanations for the matter-antimatter asymmetry in the universe.

Next-generation electric dipole moment searches may take advantage of rare isotopes such as radium-225, which are expected to be extremely sensitive to T-violation owing to their non-spherical "egg"-shaped nucleus. For the rare and unstable radium atoms, a laser trap offers a promising path to such a measurement. (Guest et al., *Phys. Rev. Lett.* **98**, 093001 (2007))

## Slowed Light Handed Off

Several years ago, physicists gained the ability to slow a beam of light in a gas of atoms; by manipulating the atoms' spins, the energy and information contained in the light could be transferred to the atoms in a coherent way. By turning on additional laser beams, the original light signal could be reconstituted and sent on its way.

Now, one of the first researchers to slow light, Lene Hau of Harvard, has added an extra layer to this story. She and her colleagues halt and store a light signal in a Bose-Einstein condensate (BEC) of sodium atoms, then transfer the signal, now in the form of a coherent pulse of atom waves rather than light waves, into a second BEC of sodium atoms some 160 microns away, from which, finally, the signal is revived as a conventional light pulse.

This feat, the sharing around of quantum information in light-form and in not just one but two atom-forms, offers great encouragement to those who hope to develop quantum computers. (Ginsberg et al., *Nature* **445**, 623-626 (8 February 2007))

## String Theory Explains RHIC Jet Suppression

String theory argues that all matter is composed of string-like shreds in a 10-dimensional hyperspace assembled in various forms. The theory has been put into play in the realm of high-energy ion collisions, the kind carried out at Brookhaven's Relativistic Heavy Ion Collider (RHIC). A few years ago string practitioners attempted to establish a relationship between the 10-dimensional string world and the 4-dimensional (3 spatial dimensions plus time) world in which we observe interactions among quark-filled particles like protons.

This duality between string theory and the theory of the strong nuclear force, quantum chromodynamics (QCD), was recently used to interpret puzzling early results from RHIC, namely the suppression of energetic quark jets that should have emerged from the fireball formed when two heavy nuclei collide head on. The thinking was that perhaps the plasma of quarks and gluons wasn't a gas of weakly interacting particles (as was originally thought) but a gas of strongly interacting particles, so strong that any energetic quarks that might have escaped the fireball (initiating a secondary avalanche, or jet of quarks) would quickly be slowed and stripped of energy on their way through the tumultuous quark-gluon plasma (QGP) environment.

Two new papers by Hong Liu and Krishna Rajagopal (MIT) and Urs Wiedemann (CERN) address this problem. The first paper calculates a specific quark-suppression parameter (namely, how much the quarks, each attached to a string dangling "downward" into a fifth dimension, are pushed around as they traverse the quark-gluon plasma) that agrees closely with the experimentally observed value.

Rajagopal says that in the second paper, the same authors make a specific testable prediction using string theory that bears not just on missing jets of energetic light quarks (up, down, and strange quarks), but on the melting or dissociation temperatures of bound states of heavy quarks (charm-anticharm or bottom-antibottom pairs) moving through the quark-gluon plasma with sufficiently high velocity, as will be produced in future experiments at RHIC and the Large Hadron Collider (LHC) under construction at CERN. (Liu, Rajagopal, and Wiedemann, *Phys. Rev. Lett.* **97**, 182301 (2006) and *Phys. Rev. Lett.* **98**, 182301 (2007))

## The Woodstock of Physics

The famous session at the 1987 March Meeting of the American Physical Society earned its nickname because of the rock-concert fervor inspired by the convergence of dozens of reports all bearing on copper-oxide superconductors. The 20th anniversary of this singular event was celebrated at the APS March Meeting in Denver.



Woodstock press conference, from left to right: Alex Muller (IBM), Paul Chu (University of Houston), Philip Anderson (Princeton), and Brian Maple (UC San Diego)

Prior to 1987 the highest temperature at which superconductivity had been observed was around 23 K. And suddenly a whole new set of compounds—not metallic alloys but crystals whose structure put them within a class of minerals known as perovskites—with superconducting transition temperatures above 35 K, and eventually 100K—generated an explosion of interest among physicists. Because of the technological benefits possibly provided by high-temperature superconductivity (HTSC)—things like bulk power storage and magnetically levitated trains—the public was intrigued too.

The commemoration of the Woodstock moment provided an excellent history lesson on how adventurous science is conducted. Georg Bednorz (IBM-Zurich), who with Alex Mueller made the initial HTSC discovery, recounted a story of frustration and exhilaration, including working for years without seeing clear evidence for superconductivity; having to use borrowed equipment after hours; overcoming skepticism from IBM colleagues and others who greatly doubted that the cuprates could support supercurrents, much less at unprecedented temperatures; and finally arriving at the definitive result—superconductivity at 35 K in a La-Ba-Cu-O compound.

In October 1986 Bednorz and Mueller prepared a journal article confirming their initial finding in the form of observing the telltale expulsion of magnetism (the Meissner effect) from the material during the transition to superconductivity. A year later Bednorz and Mueller won the Nobel Prize.

The IBM finding was soon seconded by work in Japan and at the University of Houston, where Paul Chu, testing a Y-Ba-Cu-O compound, was the first to push superconductivity above the temperature of liquid nitrogen, 77 K. Very quickly a gold rush began, with dozens of condensed matter labs around the world dropping what they were doing in order to irradiate, heat, chill, squeeze, and magnetize the new material.

At the March APS Meeting Chu said that he and his colleagues went for months on three hours' sleep per night. Several other speakers at the 2007 session spoke of the excitement of those few months in 1987 when—according to such researchers as Marvin Cohen (UC Berkeley) and Douglas Scalapino (UC Santa Barbara)—the achievement of room-temperature superconductivity did not seem inconceivable.

The Woodstock event, featuring 50 speakers delivering their fresh results at a very crowded room at the New York Hilton Hotel until 3:15 am, was a culmination. In following years, HTSC progress continued on a number of fronts, but expectations gradually became more pragmatic. Paul Chu's Y-Ba-Cu-O compound, under high-pressure conditions, still holds the transition temperature record at 164 K. Making lab samples had been easy compared to making usable power-bearing wires in long spools, partly because of the brittle nature of the ceramic compounds and partly because of the tendency for potentially superconductivity-quenching magnetic vortices to form in the material.

Paul Grant, in 1987 a scientist at IBM-Almaden, pointed out that HTSC applications have largely not materialized. No companies are making a profit from selling HTSC products. Nevertheless, the mood of the 2007 session (Woodstock20) was upbeat. Bednorz said the 1986/87 work showed that a huge leap forward could still take place in a mature research field whose origins dated back some 70 years. Bednorz felt that another wave of innovation could occur. Paul Chu ventured to predict that within ten years, HTSC products would have an impact in the power industry.

Paul Grant referred to the study of superconductivity as the “cosmology of condensed matter physics,” meaning that even after decades of scrutiny there was still much more to learn about these materials in which quantum effects, manifested over macroscopic distances, conspire to make electrical resistance vanish, a phenomenon which at some basic level might also be related to the behavior of protons inside an atomic nucleus and to the cores of distant neutron stars.

## Hyperactive Antifreeze Proteins

Hyperactive antifreeze proteins naturally secreted by an insect known as the spruce budworm prevent it from freezing to death during winters in North American forests. Ohio University's Ido Braslavsky and his colleagues presented studies of these potent yet nontoxic proteins at the APS March Meeting.

Found in several other species such as snow fleas, the hyperactive proteins bind to ice, modify its crystalline shape, and prevent ice from growing further, effectively reducing the freezing point of ice for an organism that excretes them. These nontoxic substances have more recently been renamed “ice structuring proteins” (ISPs) to distinguish them from the toxic antifreeze products for automobiles.

Extracting ISPs from biological sources has many potential applications, such as preserving organs and blood products, protecting against agricultural frost damage, and even preventing frostbite. These natural proteins are currently used in some “light” ice cream products to improve their texture, but those ISPs, derived from fish, are much less potent.

How the hyperactive versions inhibit ice from growing is a topic of interest to Braslavsky's group and their collaborators, such as Peter Davies from Queen's University. The researchers attached fluorescent molecules, derived from jellyfish, to the protein.

Through a microscope, they watched how the fluorescing ISPs inhibited ice crystals from growing. They observed that the ISPs prevent ice crystals from expanding in their normal disk-shaped form. Instead, they inhibit ice growth in certain directions and cause the crystals to grow in altered shapes.

While a fish ISP promotes the growth of a “bipyramidal” ice-crystal form that looks like two pyramids whose bases are attached to each other, the spruce budworm ISP blocks growth in the preferred direction of the pyramid's apexes. Using the fluorescence microscopy, they watched the proteins attached to the ice blocking growth in this direction. (Meeting Paper J35.8, <http://meetings.aps.org/Meeting/MAR07/Event/58982>; for more information, see <http://www.phy.ohiou.edu/~braslavs/APS2007/>)

## Quantized Magnetoresistance

The conversion of a tiny magnetic flux into a change in the resistance of an external circuit, a process called magnetoresistance, is at the heart of the \$60-billion magnetic hard-disk-drive industry. Digital data, stored on the disk in the form of minuscule domains only 50 by 200 nm in size, representing a 1 or a 0, are read out by a sensor flying only 10 nm overhead.

The first unambiguous observation of a digital version of the magnetoresistance effect—the change in the resistance recorded by the sensor changes in discrete steps as the magnetization orientation relative to the sensor is changed—was reported by physicists from the University of Nebraska and the Institut de Physique et de Chimie des Matériaux de Strasbourg (France).

The quantization of conductance on the sensor side was achieved by having the current flow through a constriction that tapers down to the size of a single atom, a passage which imposes quantum conditions. According to Nebraska scientist Andrei Sokolov, an atom-sized point contact makes the read-write process ever more compact in physical extent, allowing much greater data storage. (Sokolov et al., *Nature Nanotechnology* **2**, 171-175 (2007))

## The Ever-Shifting Face of Plutonium

A new theory explains some of the unusual properties of plutonium, the radioactive metal best known for its proclivity to undergo nuclear fission chain reactions, making it a potent fuel for nuclear weapons and power plants. Plutonium is one of the most unusual metals—it's not magnetic and it does not conduct electricity well. The material also changes its size dramatically with even the slightest changes in its temperature and pressure. The atom's unusual set of properties distinguishes it from even its closest neighbors on the periodic table, such as americium.

What makes plutonium unique? In the new theory, developed by condensed-matter theorists at Rutgers University in New Jersey, plutonium's eight outermost or “valence” electrons can circulate among different orbitals, or regions around the atom. In plutonium's 5f orbital, the one with the greatest influence on its atomic properties, the number of valence electrons it contains is most often five (approximately 80% of the time), but can also be six (about 20% of the time) or four (less than 1% of the time), according to the theory. These electrons shuttle in and out of the 5f orbital very quickly—on the order of femtoseconds, or quadrillionths of a second, the researchers say.

Plutonium is an example of a strongly correlated material, in which the valence electrons interact with each other to a great degree, and cannot be treated as independent agents. Taking these interactions into account, the researchers combined two theoretical approaches to solid materials, called the local density approximation and dynamical mean field theory, to come up with their sophisticated analysis.

As their analysis shows, the 5f orbital dictates many of plutonium's key properties, such as its lack of conductivity and magnetism. With their theory, the researchers have also explained the magnetic and electrical properties of americium and curium. They hope their approach will also elucidate the properties of rare-earth elements on the periodic table. (Shim et al., *Nature* **446**, 513-516 (29 March 2007))

## Electron Tunneling in Atoms Has Now Been Observed in Real Time

Electron tunneling in atoms has now been observed in real time by a German-Austrian-Dutch team (Ferenc Krausz, Max Planck Institute of Quantum Optics, and Ludwig Maximilians, University of Munich) using light pulses lasting only several hundred attoseconds (billionths of a billionth of a second), providing new glimpses into an important ultrafast process in nature.

The tunneling process is responsible for the operation of certain electronic components, such as scanning tunneling microscopes, Esaki (tunneling) diodes, and quantum-cascade lasers. And in nuclear fission, alpha particles are believed to escape the fracturing nucleus through tunneling. Yet the tunneling process occurs so quickly, on the scale of attoseconds, that it has not been possible to observe directly. With the recent ability to create attosecond-scale light pulses—pioneered by Krausz and others—this is now possible.

In the new experiment, a gas of neon atoms is exposed to two light pulses. One is an intense pulse containing low-energy red photons. The second pulse is an attosecond-length pulse of ultraviolet light. This ultraviolet attosecond pulse delivers photons so energetic that they can rip off an electron and promote a second one to the periphery of the atom, into an excited quantum state.

Then, the intense red pulse, consisting of just a few wave cycles, has a chance to liberate the outlying electron via light-field-induced tunneling. Indeed, the researchers saw this phenomenon, predicted theoretically forty years ago but only verified now for the first time experimentally in a direct time-resolved study. As each wave crest in the few-cycle red pulse coursed through the atoms, the electrons each time upped their probability of escaping through tunneling until it reached about 100%.

The data indicate that, in this particular system, the electrons escape via tunneling in three discrete steps, synchronized with the three most intense wave crests at the center of the few-cycle laser wave. Each step lasts less than 400 attoseconds. (Uiberacker et al, *Nature* **446**, 627-632 (5 April 2007))

## Laser Cooling of Coin-sized Objects

Laser cooling of coin-sized objects down to one-kelvin temperatures is now possible. In a set of experiments performed last year, a variation on the laser-cooling technique used in chilling vapors of gases down to sub-kelvin temperatures had been used in macroscopic (but still tiny) samples in the nano- and micro-gram range.

Now, a collaboration of scientists from the LIGO Laboratory at MIT and Caltech and from the Max Planck Institutes in Potsdam and Hanover has used laser beams to cool a coin-sized mirror with a mass of 1 gram down to a temperature of 0.8 K. The goal of chilling such a comparatively large object (with more than  $10^{20}$  atoms) is to investigate the

quantum properties of large ensembles of matter.

An important caveat here is the fact that in all these experiments the “cooling” takes place in one dimension only. A temperature of 1 K applies to the motion of atoms along the direction of the laser beams, while the mirror is free to move (although not much) in other directions. Beyond the record low temperature achieved for an object as large as 1 gram, another interesting feature of the experiment pertains to the strength of the force exerted by the laser beams. In the chosen dimension, the beams fix the mirror so steadfastly that it’s as if it were being held in place by a spring that’s stiffer than a diamond with the same dimensions as the laser beam (long and thin). According to MIT researcher Nergis Mavalvala the sample is held by a rigidity (if the laser beam were solid) characterized by a Young’s modulus (the parameter specifying stiffness) of 1.2 tera-pascals, some 20% stiffer than diamond. (Corbitt et al., *Phys. Rev. Lett.* **98**, 150802 (2007))

## Newton’s Second Law of Motion

Newton’s second law of motion has now been tested and found to be valid at the level of  $5 \times 10^{-14}$  m/sec<sup>2</sup>. This is a thousandfold improvement in precision over the best previous test, one carried out 21 years ago (*Physical Review D*, **34**, 3240, (1986)). The new test was performed by physicists at the University of Washington using a swiveling torsion pendulum.

One implication of Newton’s law is that the pendulum’s frequency should be independent of the amplitude of its swiveling (as long as the oscillation is small). Looking for a slight departure from this expected independence, the Washington researchers watched the pendulum at very small amplitudes; in fact the observed swivel was kept so small that the Brownian excitation of the pendulum was a considerable factor in interpreting the results.

Newton’s second law is expected to break down for subatomic size scales, where quantum uncertainty frustrates any precise definition of velocity. But for this experiment, where the pendulum has a mass of 70 g and consists of  $10^{24}$  atoms, quantum considerations were not important. According to one of the scientists involved, Jens Gundlach, this new affirmation that force is proportional to acceleration (at least for non-relativistic speeds), might influence further discussion of two anomalies:

(1) oddities in the rotation curves for galaxies—characterizing the velocity of stars as a function of their radii from the galactic center—suggest either that extra gravitational pull in the form of the presence of as-yet-undetected dark matter is at work or that some new form of Newton’s second law could be operating (referred to as Modified Newtonian Dynamics, or MOND); and (2) the ongoing mystery surrounding the unaccounted-for accelerations apparently characterizing the trajectory of the Pioneer spacecraft (see <http://www.aip.org/pnu/1998/split/pnu391-1.htm>). (Gundlach et al., *Phys. Rev. Lett.* **98**, 150801 (2007))

## Gravity Probe B

Gravity Probe B, the orbiting observatory devoted to testing the general theory of relativity, has measured the geodetic effect—the warping of spacetime in the vicinity of and caused by Earth—with a precision of 1%. The basic approach to studying this subtle effect is to monitor the precession of gyroscopes onboard the craft in a polar orbit around Earth. The observed precession rate, 6.6 arc-seconds per year, is close to that predicted by general relativity. Once certain unanticipated torques on the gyroscopes are better understood, GP-B scientists expect the precision of their geodetic measurement to improve to a level of 0.01%. These first GP-B results were reported at the APS April Meeting by Francis Everitt (Stanford).

A second major goal of GP-B is to measure frame dragging, a phenomenon which arises from the fact that space is, in the context of general relativity, a viscous fluid rather than the rigid scaffolding Isaac Newton took it to be. When Earth rotates, it partly takes spacetime around with it, and this imposes an additional torque on the gyroscopes.

Thus an extra precession, perpendicular to and 170 times weaker than the geodetic effect, should be observed. Everitt said that GP-B saw “glimpses” of frame dragging in this early analysis of the data and expects to report an actual detection with a precision at the 1% level by the time of the final presentation of the data.

Some of the GP-B equipment is unprecedented. The onboard telescope used to orient the gyroscopes (by sighting toward a specific star) provided a star-tracking ability better by a factor of 1000 than previous telescopes. The gyroscopes themselves—four of them for redundancy—are the most nearly spherical things ever made: the ping-pong-ball-sized objects are out of round by no more than 10 nm. They are electrostatically held in a small case and spun up to speeds of 4000 rpm by puffs of gas. The gas is then removed, creating a vacuum of  $10^{-12}$  torr. Covered with niobium and reposing at a temperature of a few kelvin, the balls are rotating superconductors, and as such they develop a tiny magnetic signature which can be read out to fix the sphere’s instantaneous orientation. (For more information see [einstein.stanford.edu](http://einstein.stanford.edu))

## One Neutrino Anomaly Has Been Resolved

One neutrino anomaly has been resolved while another has sprung up. A Fermilab experiment called MiniBooNE provides staunch new evidence for the idea that only three low-mass neutrino species exist. These results, reported at a Fermilab lecture and at the




Photo by Fermilab

A neutrino signal observed by the MiniBooNE experiment.

APS April Meeting in Jacksonville, Florida, seem to rule out two-way neutrino oscillations involving a hypothetical fourth type of low-mass neutrino.

Several experiments have previously shown that neutrinos, very light or even massless particles that only interact via gravity and the weak nuclear force, lead a schizoid life, regularly transforming from one species into another. These neutrino oscillations were presumably taking place among the three known types recognized by the standard model of particle physics: electron neutrinos, muon neutrinos, and tau neutrinos.

However, one experiment, the Liquid Scintillator Neutrino Detector (LSND) experiment at Los Alamos, provided a level of oscillation that implied the existence of a fourth neutrino species, a “sterile neutrino,” so-called because it would interact only through gravity, the weakest of physical forces.

From the start, this result stood apart from other investigations, especially since it sug-

gested possible neutrino masses very different from those inferred from the study of solar or atmospheric neutrinos or from other accelerator-based neutrino experiments. MiniBooNE set out to resolve the mystery.

The experiment proceeds as follows: protons from Fermilab’s booster accelerator are smashed into a fixed target, creating a swarm of mesons which very quickly decay into secondary particles, among them a lot of muon neutrinos. Five hundred meters away is the MiniBooNE detector. Although muon neutrinos might well oscillate into electron neutrinos, over the short run from the fixed target to the detector one would expect very few oscillations to have occurred.

The Fermilab detector, and the LSND detector before it, looked for electron neutrinos. Seeking to address directly the LSND oscillation effect, Fermilab tried to approximate the same ratio of source-detector distance to neutrino energy. This ratio sets the amount of likely oscillation.

LSND saw a small (but, they argued, statistically significant) number of electron neutrino events. MiniBooNE, after taking into account expected background events, sees none. Thus they see no oscillation and therefore no evidence for a fourth neutrino.

Actually it’s not exactly true that they see no electron neutrinos. At low neutrino energy they do see events, and this tiny subset of the data remains a mystery, to be explored in further data-taking now underway using a beam of anti-neutrinos. At the APS meeting, MiniBooNE co-spokesperson Janet Conrad (Columbia) said that the low-energy data are robust (meaning that a shortage of statistical evidence or systematic problems with the apparatus should not be major factors) and that some new physical effect cannot be ruled out.

## Tevatron’s Higgs Quest Quickens

Physicists from Fermilab’s Tevatron collider have reported their most comprehensive summary yet of physics at the highest laboratory energies. At the APS April Meeting in Jacksonville, Florida they delivered dozens of papers on a spectrum of topics, many of which are related in some way to the Higgs boson.

The Higgs is the cornerstone ingredient in the standard model of high energy physics. It is the particle manifestation of the curious mechanism that kicked in at an early moment in the life of the universe: the W and Z bosons (the carriers of the weak force) became endowed with mass while the photon (the carrier of the electromagnetic force) did not. This asymmetry makes the two forces very different in the way they operate in the universe.

Validating this grand hypothesis by actually making Higgs particles in the lab has always been a supreme reason for banging protons and antiprotons together with a combined energy of 2 TeV. However, the search for the Higgs is expected to be shadowed by the production of other rare scattering scenarios, some of them nearly as interesting as the Higgs itself.

According to Jacobo Konigsberg (University of Florida), co-spokesperson for the CDF collaboration (one of the two big detector groups operating at the Tevatron), the search for the Higgs is speeding up owing to a number of factors, including the achievement of more intense beams and increasingly sophisticated algorithms for discriminating between meaningful and mundane events.

Here is a catalog of some of the recent results from the Tevatron. Kevin Lannon (Ohio State) reported a new best figure (170.9 GeV, with an uncertainty of 1%) for the mass of the top quark. Lannon also described the class of event in which a proton-antiproton smashup resulted in the production of a single top quark via a weak-force interaction, a much rarer event topology than the one in which a top-antitop pair is made via the strong force.

Moreover, observing these single-top events allows a first rudimentary measurement of  $V_{tb}$ , a parameter proportional to the likelihood of a top quark decaying into a bottom quark. Gerald Blazey (Northern Illinois), former co-spokesperson of the D0 collaboration, reported on the first observations of equally exotic collision scenarios, those that feature the simultaneous production of an observed W and Z boson, and those in which two Z bosons are observed.

Furthermore, he said that when the new top mass is combined with the new mass for the W boson, 80.4 GeV, one calculates a new likely upper limit on the mass of the Higgs. This value, 144 GeV, is a bit lower than before, making it just that much easier to create energetically. Ulrich Heintz (Boston University) reported on the search for exotic particles not prescribed by the standard model.

Again, no major new particles were found, but further experience in handling myriad background phenomena will help prepare the way for what Tevatron scientists hope will be their main accomplishment: digging evidence for the Higgs out from a rich seam of other particles.

## The Efimov Effect: Three’s Company, Two’s a Crowd

At the APS April Meeting in Jacksonville, physicists discussed the recent observations of the Efimov effect, a purely quantum phenomenon whereby two particles such as neutral atoms which ordinarily do not interact strongly with one another join together with a third atom under the right conditions. The trio can then form an infinite number of configurations, or put another way, an infinite number of “bound states” that hold the atoms together.

The effect was first predicted around 1970 by Vitaly Efimov, then a PhD candidate, but was originally considered “too strange to be true,” according to the University of Colorado’s Chris Greene, in part because the atoms would abruptly switch from being standoffish to becoming stuck-together Siamese Triplets at remarkably long distances from one another (approximately 500-10,000 times the size of a hydrogen atom in the case of neutral atoms). For decades, experimenters tried in vain to create these three-particle systems (which came to be known as “Efimov trimers”).

In 1999, Greene and his collaborators Brett Esry and Jim Burke predicted that gases of ultracold atoms might provide the right conditions for creating the three-particle state. In 2005, a research team led by Rudi Grimm of the University of Innsbruck in Austria finally confirmed the Efimov state in an ultracold gas of cesium cooled to just 10 nanokelvin.

How do the neutral atoms attract one another in the first place? At small distances, ordinary chemical bonding mechanisms apply, but at the vast distances relevant to the Efimov effect, it is mainly through the van der Waals effect, in which rearrangements of electrical charge in one atom (forming an “electric dipole”) create electric fields that can induce dipoles in, and thereby attract, neighboring atoms.

The observation of the Efimov effect is a coup in the study of the rich quantum physics between three particles. The effect can conceivably occur in nucleons or molecules (and any object governed by quantum mechanics). However, it will likely be harder to observe in those systems because physicists cannot alter the strengths of interactions between the constituent particles as easily as they can in ultracold atom gases (through their “Feshbach

resonances”).

But the effect could provide insights on such systems as the triton, a nucleon with one proton and two neutrons, in addition to the BCS-BEC crossover, in which atoms switch from forming weakly bound Cooper pairs to entering a single collective quantum state. (See also article by Charles Day, *Physics Today*, April 2006, Esry et al., *Phys. Rev. Lett.* **83**, 1751-1754 (1999), and Kraemer et al., *Nature* **440**, 315-318 (16 March 2006).

## The Shortest Light Pulse Ever

Researchers in Italy have created the shortest light pulse yet—a single isolated burst of extreme-ultraviolet light that lasts for only 130 attoseconds (billionths of a billionth of a second). Shining this ultrashort light pulse on atoms and molecules can reveal new details of their inner workings—providing benefits to fundamental science as well as potential industrial applications such as better controlling chemical reactions.

Working at Italy’s National Laboratory for Ultrafast and Ultraintense Optical Science in Milan (as well as laboratories in Padua and Naples), the researchers believe that their current technique will allow them to create even shorter pulses well below 100 attoseconds. In previous experiments, longer pulses, in the higher hundreds of attoseconds, have been created.

The general process for this experiment is the same. An intense infrared laser strikes a jet of gas (usually argon or neon). The laser’s powerful electric fields rock the electrons back and forth, causing them to release a train of attosecond pulses consisting of high-energy photons (extreme ultraviolet in this experiment).

Creating a single isolated attosecond pulse, rather than a train of them, is more complex. To do this, the researchers employ their previously developed technique for delivering intense short (5 femtosecond) laser pulses to an argon gas target. They use additional optical techniques (including the frequency comb that was a subject of the 2005 Nobel Prize in Physics) for creating and shaping a single attosecond pulse.

The results were presented (paper JThA5) at the Conference on Lasers and Electro-Optics and the Quantum Electronics and Laser Science Conference (CLEO/QELS); also see Sansone et al., *Science* **20** October 2006: Vol. 314. no. 5798, pp. 443 - 446.)

## Ripping Fluids

A major difference between a solid and liquid is that if you move a knife through a solid, the cleft portions stay cleft, whereas in a liquid the two parts flow back together. Almost always, however, nature provides materials and processes that don’t quite fit into such neat categories.

Joseph Gladden (University of Mississippi) and Andrew Belmonte (Penn State) have contrived an experiment in which a cylinder is dragged through a mixture of water, soap, and certain salts. At small drag speeds, the material—a viscoelastic gel-like substance which is a fluid at these temperatures—does indeed close back on itself, as a liquid normally does. At higher speeds, the cylinder creates more of a cleft and the material is slower to “heal” itself. At still higher velocities, the fluid acts like a solid, at least for a while; it is ripped into several parts, with separate surfaces, which take as long as a few hours to close up, and it exhibits various “cracks” emanating from the cylinder’s wake.

Gladden says that the phase diagram (cylinder speed versus cylinder diameter) for the fluid displays three regions: flow, modest tearing, and outright ripping. Mapping out this phase diagram should help in understanding other phenomena involving viscoelastic materials, Gladden says. (*Phys. Rev. Lett.* **98**, 224501 (2007))




Image courtesy of AIP

## Nuclear Magnetic Resonance Imaging with 90-Nm Resolution

Nuclear magnetic resonance imaging with 90-nm resolution has been achieved by John Mamin and his colleagues at the IBM Almaden lab in San Jose, California. The approach used, magnetic resonance force microscopy (MRFM), maps the location of matter at small scales by observing the resonant vibration of a spindly sliver of silicon (bearing the sample in question) when it is both exposed to radio-frequency waves and scanned over a tiny magnetic tip.

Previously this same group of physicists had used a similar setup to detect the magnetic resonance of a single unpaired electron in a sample. But now they are detecting the magnetic resonance of nuclei in the sample, a much more difficult thing since nuclear magnetism is much weaker than electron magnetism (in the case of hydrogen, some 660 times weaker). The advantage in focusing on nuclear magnetism is that the response of various biologically and technologically important atoms such as H, P, C-13 or F can be differentiated.

Nuclear spin MRFM has been performed before but only with micron-scale resolution. The new imaging, in effect, explores volumes as small as 650 zeptoliters, which is some 60,000 times better than the best conventional MRI can do. Improvements in the imaging process were facilitated by the use of lower temperatures (reducing the thermally driven motion in the cantilever) and the use of very sharp magnetic tips, which enhances the magnetic force due to the spins.

The magnetic field gradient in the vicinity of this tip is greater than a million tesla/meter. The test objects being imaged consisted of tiny islands of calcium fluoride evaporated onto the cantilever tip. Closely spaced islands, roughly 300 nm x 180 nm x 80 nm in size, could be clearly resolved. One of the researchers, Dan Rugar, says that the tiny sample volumes being interrogated hold about 10 million nuclear spins, and that the net nuclear polarization they are detecting adds up to about 3300 spins.

He believes, however, that their current apparatus can now detect nuclear magnetism at the level of 200 spins. This would take them much closer to their ultimate goal of imaging




Image courtesy of AIP

molecules at the single nuclear spin level. Mamin et al., *Nature Nanotechnology* **2**, 301-306 (2007))

## Warm the World, Shrink the Day

Global warming is expected to raise ocean levels and thereby effectively shift some ocean water from currently deep areas into shallower continental shelves, including a net transfer of water mass from the southern to the northern hemisphere. This in turn will bring just so much water closer to Earth’s rotational axis, and this—like a figure skater speeding up as she folds her limbs inward—will shorten the diurnal period.

Not by much, though. According to Felix Landerer, Johann Jungclauss, and Jochem Marotzke, scientists at the Max Planck Institute for Meteorology in Hamburg, the day should shorten by 0.12 milliseconds over the next two centuries. (Landerer, Jungclauss, and Marotzke, *Geophys. Res. Lett.*, **34**, L06307 (2007))

## Microfluidic Accelerator

Microfluidics is the science of carrying out fluid chemical processing on a chip whose channels are typically millimeters or microns across. In such a constricted space, viscosity becomes large, and the fluid flow can slow way down, thus limiting the kind of mixing or testing that can be done. Physicists at the University of Twente in the Netherlands, however, use tiny exploding bubbles to speed things up.

The bubbles are produced by shooting laser light into the fluid. (See movie at [http://stilton.tnw.utwente.nl/people/ohl/controlled\\_cavitation.html](http://stilton.tnw.utwente.nl/people/ohl/controlled_cavitation.html)) The light brings a tiny volume of fluid above its boiling temperature, causing a local bubble explosion, which accelerates the surrounding fluid along the channel, now at speeds of up to 20 m/sec, twenty times higher than would be the case without the bubble, and still another factor of 10 within reach. (The same researchers have produced sonoluminescence in the same way.)

An extra advantage of using flexibly positioned laser light is that for transparent microfluidic chips, fluid pumping can be accomplished without external connections to the chip. Besides being the first to apply such a cavitation technique for speeding up fluids on a chip, the Twente scientists are the first to achieve flow visualization at rates of a million frames per second at a size scale of 100 microns.

The leader of the Twente group, Claus-Dieter Ohl, says that he and his colleagues are currently using the bubble acceleration technique for improving mixing in various enzyme reactions and in producing tiny pores in membranes. (Zwaan et al., *Phys. Rev. Lett.* **98**, 254501 (2007))

## Polonium Is the Only Element with Simple Cubic Crystal Structure

Polonium is the only element with a simple cubic crystal structure, and new theoretical work explains why that is. In a solid piece of polonium the atoms sit at the corners of a cubic unit cell and nowhere else.

One reason the study of Po is so difficult is that it is highly radioactive and spews forth decay products; indeed, polonium has more isotopes, 36, than any other element. Physicists at the Academy of Sciences in the Czech Republic have now produced the first detailed theoretical explanation for polonium’s unique crystal structure: it is the result of the complicated interplay of relativistic effects which become important in such heavy atoms as polonium (element 84).

Specifically they have identified the so-called mass-velocity term (describing the relativistic increase in mass of electrons traveling with velocities comparable to the velocity of light) as the cause of the simple-cubic structure of polonium.

Another polonium oddity: its elastic anisotropy is greater than for any other solid. That is, it is about 10 times easier to deform a Po crystal along the direction diagonal to the consolidated cubic cells than it is

to deform the crystal in a direction perpendicular to any of the cubic faces. According to Dominik Legut, this property results directly from the simple cubic structure of polonium.

Polonium is a hazardous element that appears in the air and soil and in such plants as tobacco, tea, and mushrooms. (Legut et al., *Phys. Rev. Lett.* **99**, 016402 (2007))




Image courtesy of AIP

## First Direct Measurement of DNA Stacking Forces

DNA is one of the most important and studied molecules around, and yet only now has a team of scientists, working at Duke University, succeeded in measuring the force between the nucleotides in a single-stranded DNA (ssDNA) molecule, using an atomic force microscope (AFM).

A double-stranded DNA is characterized by two principal forces—the stacking force between base units along the length of the double helix and the pairing force (Watson-Crick pairing) between the opposing base units forming the rungs of the helix. Measurements of DNA elasticity dating back to the 1990s (see <http://www.aip.org/pnu/1997/split/pnu312-1.htm>) were done with double-stranded DNA, and it is difficult to separate the effects of the pairing and stacking forces.

That’s why Piotr E. Marszalek and his colleagues (Changhong Ke, Michael Humeniuk, and Hanna S-Gracz) turned to ssDNA. They rigged an artificial ssDNA consisting only of adenine base units attached to a gold substrate, and then pulled it with an AFM tip.

With a force resolution of about 1 pico-Newton, the Duke apparatus detected one plateau in elasticity (of the stacking force) at around 23 pN, which was expected, and then a second plateau around 113 pN. (Ke et al., *Phys. Rev. Lett.* **99**, 018302 (2007)) a paper measuring forces for a single RNA molecule, finding a single force plateau at 20 pN, appeared in Seol et al., *Phys. Rev. Lett.* **98**, 158103 (2007))

## Time and Time Again

The physics world accepts the idea of spacetime, a combined metrical entity which puts

time on the same footing as the visible three spatial dimensions. Further spatial dimensions are added in some theories to help assimilate all physical forces into a unified model of reality. But what about adding an extra dimension of time too? Itzak Bars and Yueh-Cheng Kuo of the University of Southern California do exactly that, and add an extra spatial dimension too.

The addition of an extra time and an extra space dimension, together with a requirement that all motion in the enlarged space be symmetric under an interchange of position and momentum at any instant, reproduces all possible dynamics in ordinary spacetime, and brings to light many relationships and hidden symmetries that are actually present in our own universe.

The hidden relationships among dynamical systems are akin to relationships that exist between the multiple shadows of a 3D object projected on a 2D wall. In this case the object is in a spacetime of 4 space and 2 time dimensions while the shadows are in 3 space and 1 time dimensions. The motion in 4+2 dimensions is actually much more symmetric and simpler than the complex motions of the shadows in 3+1 dimensions.

In addition, Bars says that his theory explains CP conservation in the strong interactions described by QCD without the need for a new particle, the axion, which has not been found in experiments.

It also explains the fact that the elliptical orbit of planets remains fixed (not counting well-known tiny precessions). This “Runge-Lenz” symmetry effect has remained somewhat mysterious in the study of celestial mechanics, but now could be understood as being due to the symmetry of rotations into the fourth space dimension.

A similar symmetry observed in the spectrum of hydrogen would also be accounted for in 2-time physics, and again explained as a symmetry of rotations into the extra space and time dimensions. There are many such examples of hidden symmetries in the macroscopic classical world as well as in the microscopic quantum world, Bars argues, which can be addressed for the first time with the new 2T formulation of physics.

There have been previous attempts to formulate theories with a second time axis, but Bars says that most of these efforts have been compromised by problems with unitarity (the need for the sum of all probabilities of occurrences to be no greater than 1) and causality (maintaining the thermodynamic arrow of time).

The USC theorists have reformulated their model to fit into the ongoing supersymmetry version of the standard model and expect their ideas to be tested in computer simulations and in experiments yet to come. (Bars and Kuo, *Phys. Rev. Lett.* **99**, 041801 (2007))

## All-Optical Magnetic Recording

All-optical magnetic recording has been demonstrated by scientists at the Radboud University Nijmegen in the Netherlands. Instead of using the customary magnetic read head to flip the magnetic orientation of a tiny domain, they use the fields present in a short burst of circularly polarized light.

Why use light instead of a magnet? Because the magnet is relatively slow and because the magnetic field in the light pulse is intrinsically strong—up to 5 Tesla. The pulses are per-




Image courtesy of AIP

pendicularly incident on the storage medium and the helicity of the light pulse establishes whether the orientation set in the domain will be up or down, or digital terms, a 1 or a 0.

Orienting the domain (writing a bit) is accomplished partly through the light's magnetism and partly through the localized heating by the pulse, which enhances the domain's magnetic susceptibility. The bit can be reversed with light of the opposite polarization.

The light pulse is so carefully focused that it addresses only one domain at a time. The speed of the writing process is set by the duration of the laser pulse, 40 fsec, upsetting certain suggestions, made not so many years ago, that the speed of recording in optical medium could not shrink below a picosecond.

True, the size of the domain is 5 microns, which is rather large. However, one of the re-

searchers, Daniel Stanciu, says he expects the domain size to get down to about 100 nm. He believes that the all-optical approach will eventually be the way of achieving the fastest writing of data in a magnetic medium. (Stanciu et al., *Phys. Rev. Lett.* **99**, 047601 (2007))

## Hydrogen-Seven

An experiment at the GANIL facility in France is the first to make, observe, identify, and characterize the heaviest isotope yet of hydrogen, H-7, consisting of a lone proton and 6 neutrons. (An earlier experiment saw some inconclusive evidence for this state—see Korshennikov et al., *Phys. Rev. Lett.* **90**, 082501 (2003).)

All of the lighter isotopes of hydrogen have previously been seen: H-1 (ordinary hydrogen), H-2 (deuterium), H-3 (tritium), and H-4 up to H-6. Technically speaking, the H-7 state (like H-4, H-5, and H-6) is not a fully bound nucleus. It is considered a resonance since (besides being very short-lived) energy is required to force the extra neutron to adhere to the other nucleons.

In a proper nucleus energy is required to remove a neutron. In the GANIL experiment, a beam of helium-8 ions (themselves quite rare) is smashed into a carbon-12 nucleus residing in a gas of butane. In a few rare occurrences, the He-8 gives one of its protons to the C-12, producing H-7 and N-13, respectively. The H-7 flies apart almost immediately into H-3 and four separate neutrons.

Meanwhile the N-13 is observed in the active-target MAYA detector, a device much like a bubble chamber, allowing its energy and trajectory to be deduced.




Image courtesy of AIP

By taking the conservation of momentum and energy into account, the fleeting existence of the H-7 is extracted from the N-13 data. A total of seven H-7 events was observed. A rough lifetime for H-7 of less than  $10^{-21}$  seconds can be inferred. The helium-8 nucleus (2 protons plus 6 neutrons) used to make the H-7 is interesting all by itself since it is believed to consist of a nuclear core with two “halo” neutrons orbiting outside.

This radioactive species must carefully be gathered up from carbon-carbon collisions (in a separate step) and then accelerated. One of the GANIL researchers, Manuel Caamaño Fresco says that one of the chief reasons for looking at H-7 is to get a better handle on exotic nuclear matter.

The H-7 nucleus, during its brief existence, might consist of a H-3 core and plus two 2-neutron outriders, or maybe even a single 4-neutron blob outside. Larger still hydrogen isotopes, such as H-8 or H-9, might be observable. (Caamaño et al., *Phys. Rev. Lett.* **99**, 062502 (2007))

## Observing Magnetic Polarization in Single Atoms

Physicists from UC Berkeley and the Naval Research Lab have measured the spin properties of individual atoms added to a metal surface. They do this by first forming nm-sized triangular islands of cobalt on top of a copper crystal. The cobalt is ferromagnetic, which means that the spins of the cobalt atoms in the islands all line up together (half of the islands have their collective spins pointing up, while the other half point down).




Image courtesy of AIP

Additional magnetic atoms sprinkled on top of the islands (adatoms) have spins that interact magnetically with the underlying cobalt, causing the adatom spins to either align or anti-align with the underlying island spins. Thus when a small number of iron atoms (chromium atoms were also used) are dropped onto the islands they immediately become oriented (polarized) by contact with a cobalt island.

In this way isolated atoms (up to 5 nm apart) were prepared with a definite spin polarization state. Next the quantum energy levels of the magnetic adatoms were studied using the tip of a scanning tunneling microscope (STM) which itself had been magnetized.

The quantum energy levels of the iron and chromium adatoms were sampled by observing currents flowing from the adatoms into the STM tip. Current measured in this way will be larger or smaller depending on whether the spin polarization of the tip is aligned with or against the polarization of the individual magnetic adatoms being probed. The adatom energy states are seen to differ for spin-up and spin-down states, indicating that iron and chromium atoms couple magnetically to cobalt with opposite polarity.

One of the researchers, Michael Crommie of UCB, says that it is still too early to try to store data in the form of individual polarized atoms. Rather they are seeking to understand how the spin of a single atom is influenced by its environment, with an eye toward future spintronics and quantum information applications. (Yayon et al., *Phys. Rev. Lett.* **99**, 067202 (2007))

## Light-Driven Femtosecond Electricity




Image courtesy of AIP

Scientists in Canada foresee the use of electromagnetic fields of laser light for inducing and reversing tiny electrical currents along molecular wires without the use of a voltage applied across leads. They would accomplish this feat by shining special laser pulses containing light waves at two different frequencies onto a polyacetylene molecule which acts like a junction between two metallic leads on either side.

Depending on the exact frequencies used, the time duration of the pulse, and the relative phase relation

between the two components of light, the induced pulse of electric flow could consist of either a single electron or many.

For the case of one electron set in motion by the 400-femtosecond pulse of laser light, the resulting electrical “current” would be about 0.4 microamps. Why use light rather than voltage to drive electricity? Because the whole thing can be done on a femtosecond scale with lasers.

Ignacio Franco says that a potential use of laser-driven electricity would be in future optoelectronic devices such as ultrafast nanoswitches. (Franco, Shapiro and Brumer, *Phys. Rev. Lett.* **99**, 126802 (2007))

## Acoustic Quantum Dots

A new experiment at the Cavendish Lab at the University of Cambridge is the first to controllably shuttle electrons around a chip and observe their quantum properties. A quantum dot restricts electrons to a region of space in a semiconductor so tiny as to be essentially zero-dimensional. This in turn enforces a quantum regime; the electron may only have certain discrete energies, which can be useful, depending on the circumstances, for producing laser light or for use in detectors and maybe even future computers.




Image courtesy of AIP

A quantum dot is usually made not by carving the semiconductor into a tiny grain but rather by imposing restrictions on the electron's possible motions by the application of voltages to nearby electrodes. This would be a static quantum dot. It is also possible to make dynamic quantum dots—that is, moving dots that are created by the passage of surface acoustic waves (SAWs) moving through a narrow channel across the plane of a specially designed circuit chip. The acoustic wave itself is generated by applying microwaves to interleaved fingered electrodes atop a piezoelectric material like GaAs. The applied electric

fields between finger-electrodes induce a sound wave to propagate along the surface of the material.

These acoustic waves have the ability to scoop electrons and chauffeur them along the surface.

The tiny region confining the electron even as it moves is in effect a quantum dot. Such acoustic-based dynamic quantum dots have made before, but according to Cambridge researcher Michael Astley, this is the first time the tunneling of the electrons (even single electrons) into and out of the quantum dots has been observed. This is an important part of the whole electron-shuttling process since one wants control over the electron motions and spins. If, moreover, electrons in two very close acoustic wave channels could be entangled, then this would present the chance to make a sort of flying qubit, which could be at the heart of a quantum computer. (Astley et al., *Phys. Rev. Lett.* **99**, 156802 (2007))

## Thermal Logic Gates

Information processing in the world's computers is mostly carried out in compact electronic devices, which use the flow of electrons both to carry and control information. There are, however, other potential information carriers, such as photons. Indeed a major industry, photonics, has developed around the sending of messages encoded in pulsed light.

Heat pulses, or phonons, rippling through a crystal might also become a major carrier, says Baowen Li of the National University of Singapore. Li, with his colleague Lei Wang, have now shown how circuitry could use heat—energy already present in abundance in electronic devices—to carry and process information.

They suggest that thermal transistors (also proposed by Li's group in *Applied Physics Letters*, 3 April 2006) could be combined into all the types of logic gates—such as OR, AND, NOT, etc.—used in conventional processors and that therefore a thermal computer, one that manipulates heat on the microscopic level, should be possible.

Given the fact that a solid state thermal rectifier has been demonstrated experimentally in nanotubes by a group at UC Berkeley (Chang et al., *Science*, 17 November 2006) only a few years after the theoretical proposal of “thermal diode,” the heat analog of an electrical diode which would oblige heat to flow preferentially in one direction (Li et al, *Phys. Rev. Lett.* **93**, 184301 (2004)). Li is confident that thermal devices can be successfully realized in the foreseeable future. (Wang and Li, *Phys. Rev. Lett.* **99**, 177208 (2007))

## 2007 Nobel Prize in Physics




Albert Fert

The 2007 Nobel Prize in Physics was awarded to Albert Fert (Université Paris-Sud, Orsay, France) and Peter Grünberg (Forschungszentrum Jülich, Germany) for the discovery of giant magnetoresistance, or GMR for short. GMR is the process whereby a magnetic field, such as that of an oriented domain on the surface of a computer hard drive can trigger a large change in electrical resistance, thus “reading” the data vested in the magnetic orientation.

This is the heart of modern hard drive technology and makes possible the immense hard-drive data storage industry. Fert and Grünberg pioneered the making of stacks consisting of alternating thin layers of magnetic and non-

magnetic atoms needed to produce the GMR effect. GMR is a prominent example of how quantum effects (a large electrical response to a magnetic input) come about through confinement (the atomic layers being so thin); that is, atoms interact differently with each other when they are confined to a tiny volume or a thin plane.

All these magnetic interactions involve the spin of an electron. Still more innovative technology can be expected through quantum effects depending on electrons' spin. Most of the electronics industry is based on manipulating the charges of electrons moving through circuits. But the electrons' spins might also be exploited to gain new control over data storage and processing. Spintronics is the general name for this budding branch of electronics. (Nobel Prize website: [http://nobelprize.org/nobel\\_prizes/physics/laureates/2007/info.html](http://nobelprize.org/nobel_prizes/physics/laureates/2007/info.html))



Peter Grünberg

## Relativistic Thermodynamics

Einstein's special theory of relativity has formulas, called Lorentz transformations, that convert time or distance intervals from a resting frame of reference to a frame zooming by at nearly the speed of light. But how about temperature? That is, if a speeding observer, carrying her thermometer with her, tries to measure the temperature of a gas in a stationary bottle, what temperature will she measure? A new look at this contentious subject suggests that the temperature will be the same as that measured in the rest frame. In other words, moving bodies will not appear hotter or colder.

You'd think that such an issue would have been settled decades ago, but this is not the case. One problem is how to define or measure a gas temperature in the first place. James Clerk Maxwell in 1866 enunciated his famous formula predicting that the distribution of gas particle velocities would look like a Gaussian-shaped curve. But how would this curve appear to be for someone flying past? What would the equivalent average gas temperature be to this other observer? Jorn Dunkel and his colleagues at the Universität Augsburg (Germany) and the Universidad de Sevilla (Spain) could not exactly make direct measurements (no one has figured out how to maintain a contained gas at relativistic speeds in a terrestrial lab), but they performed extensive simulations of the measurement.

Dunkel says that some astrophysical systems might eventually offer a chance to experimentally judge the issue. In general the effort to marry thermodynamics with special relativity is still at an early stage. It is not exactly known how several thermodynamic parameters change at high speeds. Absolute zero, Dunkel says, will always be absolute zero, even for quickly-moving observers. But producing proper Lorentz transformations for other quantities such as entropy will be trickier to do. (Cubero et al., *Phys. Rev. Lett.* **99**, 170601 (2007))

## Nuclear Dripline Droops

Several new heavy isotopes have been discovered, at least one of which pushes beyond the neutron dripline. Driplines are the outer edges defining the zone of observed or expected bound

nuclei on a map whose horizontal axis is the number of neutrons in a nucleus (denoted by the letter N) and whose vertical axis corresponds to the number of protons (Z). The nuclear force holding neutrons and protons together (even as the like-charged protons repel each other electrostatically) is so strong that no theory (not even the so called nuclear shell model, fashioned in analogy to the atomic model) can confidently predict whether a particular combination of neutrons and protons will form a bound nucleus. Instead experimenters must help theorists by going out and finding or making each nuclide in the lab.

In an experiment conducted recently at the National Superconducting Cyclotron Lab (NSCL) at Michigan State University, a beam of calcium ions was smashed into a tungsten target. A myriad of different nuclides emerged and streamed into a sensitive detector for identification. Two newly found nuclides—Mg-40 and Al-43—came as no surprise. But another, Al-42, was more unusual since it violated the provisional prohibition against nuclei of this size having an odd number of protons and neutrons.

The new nuclides are not stable, since they decay within a few milliseconds. But this is pretty long by nuclear standards. Why study such fleeting nuclei? Even though they might not exist naturally, the new nuclides still might play a role inside stars or novae where heavy elements, including those that make up our planet and our bodies, are created. Thomas Baumann suggests that even heavier aluminum-isotopes might exist, and that it is worth exploring any possible islands of stability, not just those at the very edge of the periodic table. (Baumann et al., *Nature* **449**, 1022-1024 (25 October 2007))

## The Highest-Energy Cosmic Rays

The highest-energy cosmic rays probably come from the cores of active galactic nuclei (AGN), where supermassive black holes are thought to supply vast energy for flinging the rays across the cosmos. This is the conclusion reached by scientists who operate the Pierre Auger Observatory in Argentina. This gigantic array of detectors spread across 3000 sq. km of terrain, looks for one thing: cosmic ray showers.

These arise when extremely energetic particles strike our atmosphere, spawning a gush of secondary particles. Many of the rays come from inside our own Milky Way, especially from our sun, but many others come from far away. Of most interest are the highest-energy showers, with energies above  $10^{19}$  electron volts, far higher than any particle energy that can be produced in terrestrial accelerators. The origin of such potent physical artifacts offers physicists a tool for studying the most violent events in the universe.

To arrive at Earth, most cosmic rays will have crossed a great deal of intergalactic space, where magnetic fields can deflect them from their starting trajectories. But for the highest-energy rays, the magnetic fields can't exert as much influence, and consequently the starting point for the cosmic rays can be traced with some confidence.

This allowed the Auger scientists to assert that the highest-energy cosmic rays were not coming uniformly from all directions but rather preferentially from galaxies with active cores, where the engine for particle acceleration was probably black holes of enormous size. The very largest of cosmic ray showers, those with an energy higher than 57 EeV (1 EeV equals  $10^{18}$  eV), correlated pretty well with known AGN's. (Auger collaboration, *Science* **9** November 2007: Vol. 318, no. 5852, pp. 938-943)

## Cooper Pairs in Insulators

Cooper pairs are the extraordinary link-up of like-charged electrons through the subtle flexings of a crystal. They act as the backbone of the superconducting phenomenon, but have also now been observed in a material that is not only non-superconducting but actually an insulator. An experiment at Brown University measures electrical resistance in a Swiss-cheese-like plank of bismuth atoms made by spritzing a cloud of atoms onto a substrate with 27-nm-wide holes spaced 100 nm apart. Bismuth films made this way are superconducting if the sample is many atom-layers thick but is insulating if the film is only a few atoms thick, owing to subtle effects which arise from the restrictive geometry.

Cooper pairs are certainly present in the superconducting sample; they team up to form a non-resistive supercurrent. But how do the researchers know that pairs are present in the insulator too? By seeing what happens to resistance as an external magnetic field is increased.

The resistance should vary periodically, with a period proportional to the charge of the electrical objects in question. From the periodicity, proportional in this case to two times the charge of the electron, the Brown physicists could deduce that they were seeing doubly-charged objects moving through the sample. In other words, Cooper pairs are present in the insulator. This is true only at the lowest temperatures. One of the researchers, James Valles, says that there have been previous hints of Cooper pairs in some films related to superconductors, but that in those cases the evidence for pairs in the insulating state was ambiguous. He asserts that the realization of a boson insulator (in which the charge carriers are electron pairs) will help to further explore the odd kinship between insulators and superconductors. (Stewart et al., *Science* **23** November 2007: Vol. 318, no. 5854, pp. 1273-1275)

## Persistent Flow or Bose-Condensed Atoms in a Toroidal Trap

A persistent flow of Bose-condensed atoms has been achieved for the first time, offering physicists a better chance to study the kinship between Bose-Einstein condensates (BEC) and superfluids. Both involve the establishment of an ensemble in which many atoms join together in a single quantum entity. But they're not quite the same thing. In a bath of liquid helium at low temperatures, for example, nearly 100% of the atoms are in a superfluid state but only about 10% are in a BEC state (in a BEC millions of atoms have become, in a sense, a single atom). But physicists generally believe that most or all of a BEC is superfluid. Scientists have been able to stir up quantized vortices in BEC samples, one indication that BECs are superfluid. But until now researchers had not been able to get BECs to move around a track in a persistent flow, another sign of superfluidity.

The new experiment, performed by Nobel laureate William Phillips and his colleagues at NIST-Gaithersburg, the Joint Quantum Institute of NIST and the University of Maryland, chilled sodium atoms in a toroidal trap, set them into motion with laser light, and observed a flow for as long as 10 seconds.

One of the scientists on the project, Kristian Helmerson, says that neutral atoms flowing in a toroidal vessel could be fashioned into the atom analog of a superconducting quantum interference device (SQUID), which is used as a sensitive detector of magnetism. This BEC device, sensing not magnetism but slight changes in direction, could serve as a sensitive gyroscope, possibly for navigation purposes. (Ryu et al., *Phys. Rev. Lett.* **99**, 260401 (2007))

## Senior Physicists Group 10 Years Old and Going Strong

By Calla Cofield

In 1996 Dick Strombotne retired from a long and busy career with the U.S. Department of Transportation. Like many physicists, Strombotne may have been retired, but his love of physics and his desire to learn hadn't waned. Having been a member of APS for 40 years, Strombotne contacted APS Executive Officer Judy Franz to see if there was a local group of physicists with a similar desire to remain active in the physics community. Since there wasn't one, Strombotne started it himself. More than 50 physicists came out for the initial planning meeting, and in the past ten years, the contact list of members from the greater Washington DC area has grown to 160.

The Mid-Atlantic Senior Physicists Group is now celebrating its ten-year anniversary. For the past

members and some aren't. Some members are senior in age, some are senior in experience, and some are both, but despite the name the activities are open to anyone.

The group's featured talks have been about nanotechnology, global climate change, topics in astrophysics, string theory, supersymmetry, medical physics, and the evolution of standard time, to name a few. They usually take place on the third Wednesday of each month at APS headquarters in College Park, MD. Other activities have included a visit to the M.C. Escher exhibit at the National Gallery in Washington, and tours to different physics installations such as the David Taylor Model Basin at the Naval Research Laboratory, the Applied Physics Lab at Johns Hopkins, and a two-day trip to the National Radio Astronomy Observatory in Green



Photo by Calla Cofield

Richard Strombotne (far left) and other members of the Mid-Atlantic Senior Physicists Group listen to a talk by David Newell of NIST at the American Center for Physics in College Park, MD.

decade the group has sponsored talks on physics, tours to local physics installations, and trips to more distant sites. The events are open to anyone interested in attending.

Strombotne says of the formation of the group, "For the first year or so, we had about three talks in the spring and three talks in the fall. Now we have a talk in most months...and tours of some installation in other months." Four of the original ten planning committee members are still on board with the group.

Strombotne says they called the group "Senior Physicists" because "senior" seemed to describe the people most interested in participating. Many of the group's members are retired, but many are still working full or part time. Some are APS

Bank, West Virginia.

For the past ten years the group has existed as an informal part of APS. They are considering trying to become an APS Forum, which would require 200 members who belong to APS. As a forum, the group would gain additional funding and exposure, potentially uniting them with other regional groups of the same nature, or motivating others to start these groups where there are none. Mainly, the group would like to make themselves known to more physicists who would be interested in joining. A website for the group is scheduled for launch later in 2008, and until then anyone interested in attending a talk or activity can contact APS Director of Membership Trish Lettieri at [lettieri@aps.org](mailto:lettieri@aps.org) or 301-209-3272.

## New Lab Association Elects Officers

The Advanced Laboratory Physics Association (ALPhA) elected its first officers in November. The organization formed last spring after holding an organizational session at the 2007 APS March Meeting (see *APS News*, May 2007), and now has over 100 members.

The mission of ALPhA is to encourage communication among those who teach advanced undergraduate laboratory courses and provide professional recognition for those instructors. "ALPhA hopes to become the central advocacy group for advanced experimental physics instruction," its constitution states.

ALPhA will work closely with both the American Association of

Physics Teachers (AAPT) and APS. ALPhA will plan sessions at AAPT and APS meetings, suggest invited talks at these meetings, award professional prizes, and plan special conferences on advanced experimental instruction. The organization held an open meeting at the January AAPT meeting, and will have a similar event at the 2008 APS March Meeting.

Recently elected ALPhA president Gabe Spalding of Illinois Wesleyan University has been teaching advanced lab for 12 years, and regards these courses as challenging. "There should be thought given to the experimental curriculum," he said.

In addition to fitting in with the rest of the curriculum and teaching

appropriate experimental skills, these courses have to teach a "mindset," said Spalding. A good advanced lab course encourages students to take more ownership of their projects, he added.

"I believe that ALPhA will have an enormous impact on the instruction of the experimental curriculum. This sort of banding together of dedicated instructors can be transformative," said Spalding.

AAPT has established a listserv devoted to the teaching of advanced laboratory courses. The listserv discusses everything from technical problems with equipment to issues of curriculum, and is open to anyone. It can be found

at <http://lists.aapt.org/cgi-bin/lyris.pl?enter=advlabs-l>

Joining ALPhA costs \$10. To join, visit <http://www.teachspin.com/signup.shtml>

ALPhA's president, vice-president, and board members will serve two-year terms, the secretary and treasurer four-year terms, and the vice-president will automatically serve a second two-year term as president. The recently elected officers, most of whom are also members of APS, are:

President: **Gabriel Spalding**,  
*Illinois Wesleyan University*

Vice-President: **James Lockhart**,  
*San Francisco State University*

Now Appearing in RMP:  
Recently Posted Reviews and Colloquia  
You will find the following in the  
online edition of  
*Reviews of Modern Physics* at  
<http://rmp.aps.org>

**Four-body methods for high-energy ion-atom collisions**  
Dževad Belkić, Ivan Mančev  
and Jocelyn Hanssen

This review presents a thorough overview of the current status and a critical assessment of the existing quantum-mechanical four-body theories for energetic ion atom collisions. A proper description of these collisions with two active electrons requires the solution of the pertinent four-body problems. The analysis considers a number of inelastic collisions with special attention focused on single electron capture, double electron capture, transfer ionization, and transfer excitation. Working within the four-body framework of scattering theory and imposing proper Coulomb boundary conditions in the entrance and exit channels, the review analyzes a number of the leading quantum-mechanical theories. The scope of the review is limited to intermediate and high nonrelativistic impact energies.

Looking for a new job  
Looking to hire qualified candidates?

### APS March Meeting Job Fair

March 10-11, 2008  
New Orleans, LA

Specializing in all areas of:

- Physics
- Engineering
- Related Physical Sciences

For more information:

<http://www.aps.org/careers/employment/jobfairs.cfm>

For additional information contact:

Alix Brice at [abrice@aip.org](mailto:abrice@aip.org) or (301) 209-3187

POPA continued from page 1

hydrogen-powered vehicles competitive. The report made headlines in several major newspapers, on National Public Radio, and prompted Congress to hold hearings on the issues that were raised in the report (see *APS News*, May 2004, available online). In the end, Congress implemented a program consistent with the recommendations in POPA's short report.

In May 2005, POPA released a second short report on nuclear power and proliferation resistance titled "Securing Benefits, Limiting Risk," examining technological steps that could be taken to guard nuclear power systems from the threat of theft or diversion. Its impact can be seen in the controversial fiscal year 2008 omnibus spending bill recently passed by Congress (see story, page 1), which eliminated all funding for both the proposed nuclear fuel recycling facility and advanced burner reactor, restricting the U.S. Department of Energy's recycling efforts to a research program.

"We've been lobbying for these outcomes for the last 18 months, and it's good to see such clear and unambiguous results," says Slakey.

Up next are two short reports, to be released this month, one on nuclear workforce and the other on nuclear forensics. The latter examines the potential for nuclear forensics to enhance global nuclear deterrence. The group's charge reads, in part, "In a world with many nuclear weapons states, the ability of the US to use scientific techniques to identify with certainty the origin of the fissile material used in an attack is increasingly important."



## Shoot-Out at the O.K. Corral

By Michael S. Lubell, APS Director of Public Affairs

It was Washington, not Tombstone, and it was 2007, not 1881, but when Congress finally passed the Omnibus spending bill for Fiscal Year 2008 shortly before Christmas, their labor left many government activities riddled with holes.

Like the shoot-out at the O.K. Corral a century earlier, some of the facts surrounding the fiscal casualties still remain murky. And just as Tom McClaurry and Billy Clanton might have escaped the hail of bullets that day had Wyatt Earp and "Doc" Holliday known they were unarmed, science might have come through the partisan fusillade last year had Congress known the consequences of its actions.

But McClaurry and Clanton were mortally wounded. And though science is still alive, it is badly in need of life support.

The dispute between Earp and the "Cowboys" dated back more than a year before the shoot-out. And so, too, with the Fiscal Year 2008 science budget: the seeds of the antagonism that led to its collapse were sown in 2006.

In his State of the Union Ad-

dress that year, President Bush unveiled the American Competitiveness Initiative that put the research budgets of the Department of Energy's Office of Science, the National Institute of Standards and Technology, and the National Science Foundation on a ten-year doubling path. The President's announcement followed on the heels of the National Academies' report, *Rising Above the Gathering Storm*, and the House Democrats' release of their *Innovation Agenda*.

With Congress still in the hands of the Republicans, who had catered to the White House during the previous three years, Washington assumed that 2006 would continue the pattern. But by the time the November elections rolled around, the Republicans had failed to pass any domestic appropriations bills. And after their defeat at the polls, the GOP packed its bags, leaving behind a huge fiscal mess for the 110th Democratic Congress to clean up.

As the Republicans fled, Democrats warned they would have less than a month to deal with Fiscal Year 2007 spending before the 2008 budget landed on their desks.

**BELTWAY continued on page 12**

Secretary: **Mark Masters**,  
*Indiana University-Purdue University, Fort Wayne*

Treasurer: **Steve Wonnell**,  
*Johns Hopkins University*

Elected members of the executive board are:

**Robert DeSerio**,  
*University of Florida*

**Paul Dolan**,  
*Northeastern Illinois University*

**Richard Peterson**,  
*Bethel University*

**Jonathan Reichert**,  
*TeachSpin, Inc.*

# The Back Page

## A Physicist for President?

By Vernon J. Ehlers

The current presidential campaign cycle started earlier than usual, giving candidates abundant time to talk about the issues. Unfortunately, I have heard little discussion about science and technology. Candidates carefully avoid topics that might make them look too “nerdy,” that is, overly interested in nuclei, Euclidian geometry or theoretical chemistry.

Later this year, millions of Americans will cast their ballot to elect the next President of the United States, but few will investigate the candidates’ understanding of science. Do you know where your favorite candidate stands on science and technology issues? Have you heard any candidate explain the importance of science and math education to our national defense, energy solutions, global competitiveness, health care, or the ability of our students to obtain meaningful employment in the future? Have any discussed the necessity of adequately funding scientific research?

Before you cast your ballot, consider one additional, hypothetical candidate: “Physicist for President.” Let me first make it clear: I have absolutely no desire to run for president. But as a physicist, I hope that in my lifetime someone who holds an advanced degree in physics, or some other science, will run for and win our nation’s top office.

The physicist’s presidential platform would give science and technology prominence. The candidate would recognize that geographic boundaries are almost meaningless in the 21st century. He or she would recognize that the Internet and other technologies have allowed financial and intellectual capital to flow freely worldwide at nearly the speed of light. The United States is no longer competing with a handful of developed countries, but with the entire world.

On October 4, 2007, we recalled the 50th anniversary of the launch of Sputnik I into orbit. People who were alive in 1957 vividly remember this event. It shocked the American public and dwarfed the achievements of our rocket program. Sputnik spurred U.S. investment in aerospace, culminating in the Apollo moon landing. It also stimulated a great emphasis on improving our math and science education programs and sparked an intense focus on equipping our workforce with the skills needed to compete with the Russians and other foreign countries.

Today, the United States is facing an equally critical challenge from overseas. Despite lacking the same public prominence as the Sputnik launch, our children are once again falling behind their peers in European and Asian countries in the subjects of math and science. As a physicist, it is clear to me how important these subjects are in preparing students for the jobs of the future. I am concerned that by the time another Sputnik-like spark comes along to wake us up to the crisis looming over our nation’s competitiveness, it may be too late to act. In order to address this growing challenge, a physicist would support updating the No Child Left Behind Act, which has helped countless students in the United States improve over the past five years. This would help ensure that students are prepared for the jobs of the future.

A scientist in the Oval Office would bring good analytical skills to decision-making in the White House, and would appreciate the need for a population well-versed in science. A public which understands basic scientific principles and concepts would produce analytical voters and ensure we are better stewards of our planet and all that it contains.

A physicist’s platform would also include sustained investment in fundamental research. President Bush recently signed into law the America COMPETES Act of 2007. This law includes provisions to encourage innovation in manufacturing and to strengthen many of our federal research and education programs. It also provides incentives to increase the number of science, technology, engineering and mathematics (STEM) majors and teachers. Also, through its spe-



cial focus on the training of teachers, it seeks to improve STEM education for all of our nation’s children, not just the ones who will pursue advanced degrees. It strives to equip all high school graduates with a strong education in science and math, allowing them to excel in any career path they choose. The law establishes a pathway to double in seven years the research budgets of the National Science Foundation, the Department of Energy Office of Science, and the National Institute of Standards and Technology, and it enhances programs designed to improve K-12 teacher content knowledge in science and math. The successful passage of this \$33.6 billion authorization was in large part due to advocacy by individual scientists across the nation. I was excited to see the efforts of 12 years of hard work in Congress pay off when this bill became law, and I am pleased that I was able to play a part in this success. Clearly, with a scientist as president, as well as more scientists in Congress, success could be achieved much more rapidly.

The catch, however, is that the COMPETES Act does not ensure that this funding actually materializes, since authorization measures must be followed by Appropriation Committee actions to ensure the funds are allocated and spent. Of course, the “physicist-for-president” platform would include a plan to fully fund the COMPETES Act in the annual budget request to Congress. Though our nation’s president should have a fiscally conservative view on government spending, this would be the only part of a platform where a scientist could clearly make the case that this investment is one which we must not underfund. It is truly an investment in our future and would produce a great return on investment. Americans must recognize how important basic research is to the vitality of our nation; fully funding these programs should be a proposal all would support. Additionally, I expect a scientist running for president would pledge to permanently extend the research and development tax credit; this would give companies the ability to depend on that credit when they conduct long-term planning for their research and development endeavors.

In summary, the “Physicist for President” platform would present our nation with a winning array of ideas developed to put us on the path toward sustained economic competitiveness and bolstered innovation. It would include substan-

tial investments in our nation’s research and development programs, as well as sustained efforts to build upon our successful STEM education programs. China and India decided 20 years ago to improve the STEM education of their students, and today are reaping tangible results, especially in manufacturing. It is time for us to catch up to the substantial investments other nations are making.

Perhaps someday we will elect a scientist as president. Until then, I urge my fellow physicists to become involved in their communities and local politics. Volunteer to speak at your local high school, so you can excite students about science. Run for your local school board. Serve as a volunteer advisor to an elected official. Mentor a student and encourage him or her to pursue a college degree or career in science. Exercise your right to vote. If you interact with a real presidential candidate, ask him or her for positions on these issues. If you would like to establish a personal relationship with a candidate, I encourage you to attend their events and to volunteer to work for their campaign. Similarly, I urge current policy-

makers to listen to the voices of physicists. Our unique training provides us with the perspective to approach problems logically while analytically developing solutions.

I sincerely hope that our next President will share the same passion and zeal I have for improving our nation’s science and education programs, and that our country will grow and prosper from scientific knowledge! Wouldn’t you vote for that? And won’t you work to make it happen?

APS Fellow Vernon J. Ehlers received his PhD in nuclear physics from UC Berkeley. A Republican, he has represented Michigan’s 3rd district in Congress since 1993. He serves on the Education and Labor, Science and Technology, Transportation and Infrastructure, and Administration committees of the House of Representatives.

**BELTWAY continued from page 11**

All they could do, they said, was pass a Continuing Resolution, freezing all programs at the previous year’s level. The President’s competitive initiative, ACI, would have to be put on hold for a year.

But following an intense lobbying effort, science advocates managed to get a rare limited waiver for the three ACI agencies, and funding rose, though not as much as the President had requested. With White House science advisor Jack Marburger still sidelined by illness, the Office of Science and Technology Policy scolded the Democrats for funding only half the President’s request: not a word about the failures of the 109th Republican Congress.

The partisan attack did not sit well with Democrats. Just hours later, House Science and Technology Committee Chairman Bart Gordon (D-6th TN) responded, “While the President’s [FY 2008] budget includes some important funding increases, it lacks the priorities and consistency to ensure our competitiveness now and in the long run.”

So much for science bipartisanship! Still, in the coming months, Congress managed to pass overwhelmingly the America COMPETES Act that authorized the ACI doubling, and despite some grumbling, the President signed the bill. The House also passed spending bills with the ACI increases included. So too, did the Senate Appropriations Committee.

But the Senate leadership failed to bring any of its bills to the floor for a vote. Ultimately, the only option was an eleventh-hour \$933 billion Omnibus bill, \$22 billion above the White House bottom line.

The President held to his number, adamantly refusing even to meet with the Democrats. And in return Democrats swore that they would “whack GOP priorities” to meet his demand. They did. In the process, the ACI increases evaporated.

Was science simply caught in the crossfire, or was ACI a target? We may never know.

**“In summary, the ‘Physicist for President’ platform would present our nation with a winning array of ideas developed to put us on the path toward sustained economic competitiveness and bolstered innovation.”**

**“A scientist in the Oval Office would bring good analytical skills to decision-making in the White House, and would appreciate the need for a population well-versed in science.”**

**“Have you heard any candidate explain the importance of science and math education to our national defense, energy solutions, global competitiveness, health care, or the ability of our students to obtain meaningful employment in the future?”**